arXiv:1304.6248v2 [hep-ph] 1 May 2013

Lepton-flavor violation and $(g-2)_\mu$ in the $\mu\nu$SSM

Hai-Bin Zhang[a,b,*], Tai-Fu Feng[a,b], Shu-Min Zhao[a], Tie-Jun Gao[a,b]

[a] *Department of Physics, Hebei University, Baoding 071002, China*
[b] *Department of Physics, Dalian University of Technology, Dalian 116024, China*

Abstract

Within framework of the μ from ν Supersymmetric Standard Model ($\mu\nu$SSM), exotic singlet right-handed neutrino superfields induce new sources for lepton-flavor violation. In this work, we investigate some lepton-flavor violating processes in detail in the $\mu\nu$SSM. The numerical results indicate that the branching ratios for lepton-flavor violating processes $\mu \to e\gamma$, $\tau \to \mu\gamma$ and $\mu \to 3e$ can reach 10^{-12} when $\tan\beta$ is large enough, which can be detected in near future. We also discuss the constraint on the relevant parameter space of the model from the muon anomalous magnetic dipole moment. In addition, from the scalars for the $\mu\nu$SSM we strictly separate the Goldstone bosons, which disappear in the physical gauge.

Keywords: Lepton-flavor violation, R-parity violation, supersymmetry, anomalous magnetic dipole moment.
2010 MSC: 81T60, 81V15

1. Introduction

It is obviously evidence of new physics beyond the Standard Model (SM) that if we observe lepton-flavor violating (LFV) processes in future experiments, because the lepton-flavor number is conserved in the Standard Model. In supersymmetric (SUSY) extensions of the SM, the R-parity of a particle is defined as $R = (-1)^{L+3B+2S}$ [1] and can be violated if either the baryon number (B) or lepton number (L) is not conserved [2, 3], where S denotes

---

[*] Corresponding author

Email addresses: `hbzhang@mail.dlut.edu.cn` (Hai-Bin Zhang), `fengtf@hbu.edu.cn` (Tai-Fu Feng)

the spin of concerned component field. Note that $R = +1$ for particles and -1 for superparticles.

Differing from the models in Refs.[2, 3], the authors of Ref.[4] propose a supersymmetric extension of the SM named as the "μ from ν Supersymmetric Standard Model" ($\mu\nu$SSM), which solves the μ problem [5] of the Minimal Supersymmetric Standard Model (MSSM) [6] through the lepton number and R-parity breaking couplings between the right-handed neutrino superfields and the Higgses $\epsilon_{ab}\lambda_i\hat{\nu}_i^c\hat{H}_d^a\hat{H}_u^b$ in the superpotential. The effective μ term $\epsilon_{ab}\mu\hat{H}_d^a\hat{H}_u^b$ is generated spontaneously through right-handed sneutrino vacuum expectation values (VEVs), $\mu = \lambda_i \langle\tilde{\nu}_i^c\rangle$, as the electroweak symmetry is broken (EWSB). Note that a popular model is the so-called Bilinear R-parity Violation (BRpV) model [3], where the BRpV terms $\epsilon_{ab}\varepsilon_i\hat{H}_u^b\hat{L}_i^a$ are added to the MSSM. The effective BRpV terms are generated spontaneously through the R-parity conserved terms $\epsilon_{ab}Y_{\nu_{ij}}\hat{\nu}_j^c\hat{H}_u^b\hat{L}_i^a$ in the superpotential of the $\mu\nu$SSM, and $\varepsilon_i = Y_{\nu_{ij}}\left\langle\tilde{\nu}_j^c\right\rangle$, as EWSB. So largely differing from the other models [2, 3], the $\mu\nu$SSM introduces three exotic right-handed sneutrinos $\hat{\nu}_i^c$, and once EWSB the right-handed sneutrinos give nonzero VEVs. In addition, the nonzero VEVs of right-handed sneutrinos induce new sources for lepton-flavor violation. In this work, we analyze the constraints on parameter space of this model from the experimental observations on some LFV processes and muon anomalous magnetic dipole moment (MDM).

If the left-handed scalar neutrinos acquire nonzero vacuum expectation values when the electroweak symmetry is broken , the tiny neutrino masses are aroused [7] to account for the experimental data on neutrino oscillations [8, 9, 10]. Three flavor neutrinos $\nu_{e,\mu,\tau}$ are mixed into three massive neutrinos $\nu_{1,2,3}$ during their flight, and the mixings are described by the Pontecorvo-Maki-Nakagawa-Sakata unitary matrix U_{PMNS} [11]. The experimental observations of the parameters in U_{PMNS} for the normal mass hierarchy [12] show that [13]

$$\begin{aligned}&\Delta m_{21}^2 = 7.58^{+0.22}_{-0.26} \times 10^{-5}\text{eV}^2\ , \qquad \Delta m_{32}^2 = 2.35^{+0.12}_{-0.09} \times 10^{-3}\text{eV}^2\ ,\\ &\sin^2\theta_{12} = 0.306^{+0.018}_{-0.015}, \quad \sin^2\theta_{23} = 0.42^{+0.08}_{-0.03}, \quad \sin^2\theta_{13} = 0.021^{+0.007}_{-0.008}. \end{aligned} \tag{1}$$

Note that the Daya Bay Reactor Neutrino Experiment has measured a nonzero value for the neutrino mixing angle θ_{13} with a significance of 5.2 standard deviations recently [14]. Differing from the BRpV model, where one neutrino mass is generated at tree level and the other two at one loop [15], the $\mu\nu$SSM can generate three neutrino masses at the tree level through the mixing with the neutralinos including three right-handed neutrinos [16, 17]. Here, we use the neutrino experimental data presented in Eq.(1) to restrain the input parameters in the model. Then, we analyze the branching ratios for the various LFV processes: $\mu \to e\gamma$, $\tau \to \mu\gamma$, $\mu \to 3e$, etc., and the corrections to the anomalous magnetic dipole moment of the muon a_μ in the $\mu\nu$SSM. The numerical results indicate that the new physics contributes large corrections to the branching ratios of the mentioned LFV processes and a_μ in some parameter space of the model.

The outline of the paper is as follow. In section 2, we present the ingredients of the $\mu\nu$SSM by introducing its superpotential and the general soft SUSY-breaking terms, in particular we strictly separate the unphysical Goldstone bosons from the scalars. In section 3, we analyze the decay width of those interested rare LFV processes, and present the SUSY contribution to muon MDM in section 4. The numerical analysis is given in section 5, and the conclusions are summarized in section 6. The tedious formulae are collected in Appendices.

2. The $\mu\nu$SSM

Besides the superfields of the MSSM, the $\mu\nu$SSM introduces three exotic gauge singlet neutrino superfields $\hat{\nu}_i^c$. The corresponding superpotential of the $\mu\nu$SSM is given as [4]

$$\begin{aligned} W &= \epsilon_{ab}\Big(Y_{u_{ij}}\hat{H}_u^b\hat{Q}_i^a\hat{u}_j^c + Y_{d_{ij}}\hat{H}_d^a\hat{Q}_i^b\hat{d}_j^c + Y_{e_{ij}}\hat{H}_d^a\hat{L}_i^b\hat{e}_j^c + Y_{\nu_{ij}}\hat{H}_u^b\hat{L}_i^a\hat{\nu}_j^c\Big) \\ &\quad - \epsilon_{ab}\lambda_i\hat{\nu}_i^c\hat{H}_d^a\hat{H}_u^b + \frac{1}{3}\kappa_{ijk}\hat{\nu}_i^c\hat{\nu}_j^c\hat{\nu}_k^c\,, \end{aligned} \tag{2}$$

where $\hat{H}_d^T = \Big(\hat{H}_d^0, \hat{H}_d^-\Big)$, $\hat{H}_u^T = \Big(\hat{H}_u^+, \hat{H}_u^0\Big)$, $\hat{Q}_i^T = \Big(\hat{u}_i, \hat{d}_i\Big)$, $\hat{L}_i^T = \Big(\hat{\nu}_i, \hat{e}_i\Big)$ (the index T denotes the transposition) are $SU(2)$ doublet superfields, and $\hat{d}_j^c$, $\hat{u}_j^c$ and $\hat{e}_j^c$ represent the singlet down-type quark, up-type quark and lepton superfields, respectively. In addition, $Y_{u,d,e,\nu}$, λ and κ are dimensionless matrices, a vector and a totally symmetric tensor. a, b are SU(2) indices with antisymmetric tensor $\epsilon_{12} = -\epsilon_{21} = 1$, and $i, j = 1,\ 2,\ 3$. The summation convention is implied on repeated indices.

In the superpotential, the first three terms are almost the same as the MSSM. Next two terms can generate the effective bilinear terms $\epsilon_{ab}\varepsilon_i\hat{H}_u^b\hat{L}_i^a$, $\epsilon_{ab}\mu\hat{H}_d^a\hat{H}_u^b$, and $\varepsilon_i = Y_{\nu_{ij}}\left\langle\tilde{\nu}_j^c\right\rangle$, $\mu = \lambda_i\left\langle\tilde{\nu}_i^c\right\rangle$, once the electroweak symmetry is broken. The last term can generate the effective Majorana masses for neutrinos at the electroweak scale. And the last two terms explicitly violate lepton number and R-parity.

The general soft SUSY-breaking terms in the $\mu\nu$SSM are given by

$$\begin{aligned}
-\mathcal{L}_{soft} \;=\; & m^2_{\tilde{Q}_{ij}}\tilde{Q}^{a*}_i\tilde{Q}^a_j + m^2_{\tilde{u}^c_{ij}}\tilde{u}^{c*}_i\tilde{u}^c_j + m^2_{\tilde{d}^c_{ij}}\tilde{d}^{c*}_i\tilde{d}^c_j + m^2_{\tilde{L}_{ij}}\tilde{L}^{a*}_i\tilde{L}^a_j \\
& + m^2_{\tilde{e}^c_{ij}}\tilde{e}^{c*}_i\tilde{e}^c_j + m^2_{H_d}H^{a*}_dH^a_d + m^2_{H_u}H^{a*}_uH^a_u + m^2_{\tilde{\nu}^c_{ij}}\tilde{\nu}^{c*}_i\tilde{\nu}^c_j \\
& + \epsilon_{ab}\Big[(A_uY_u)_{ij}H^b_u\tilde{Q}^a_i\tilde{u}^c_j + (A_dY_d)_{ij}H^a_d\tilde{Q}^b_i\tilde{d}^c_j + (A_eY_e)_{ij}H^a_d\tilde{L}^b_i\tilde{e}^c_j + \text{H.c.}\Big] \\
& + \Big[\epsilon_{ab}(A_\nu Y_\nu)_{ij}H^b_u\tilde{L}^a_i\tilde{\nu}^c_j - \epsilon_{ab}(A_\lambda\lambda)_i\tilde{\nu}^c_iH^a_dH^b_u + \frac{1}{3}(A_\kappa\kappa)_{ijk}\tilde{\nu}^c_i\tilde{\nu}^c_j\tilde{\nu}^c_k + \text{H.c.}\Big] \\
& - \frac{1}{2}\Big(M_3\tilde{\lambda}_3\tilde{\lambda}_3 + M_2\tilde{\lambda}_2\tilde{\lambda}_2 + M_1\tilde{\lambda}_1\tilde{\lambda}_1 + \text{H.c.}\Big)\,. \qquad (3)
\end{aligned}$$

Here, the front two lines contain squared-mass terms of squarks, sleptons and Higgses. The next two lines consist of the trilinear scalar couplings. In the last line, M_3, M_2 and M_1 denote Majorana masses corresponding to $SU(3)$, $SU(2)$ and $U(1)$ gauginos $\hat{\lambda}_3$, $\hat{\lambda}_2$ and $\hat{\lambda}_1$, respectively. In addition to the terms from $\mathcal{L}_{soft}$, the tree-level scalar potential receives the usual D and F term contributions [4].

When the electroweak symmetry is spontaneously broken (EWSB), the neutral scalars develop in general the vacuum expectation values (VEVs):

$$\langle H^0_d\rangle = \upsilon_d, \qquad \langle H^0_u\rangle = \upsilon_u, \qquad \langle\tilde{\nu}_i\rangle = \upsilon_{\nu_i}, \qquad \langle\tilde{\nu}^c_i\rangle = \upsilon_{\nu^c_i}. \qquad (4)$$

Thus one can define neutral scalars as usual

$$\begin{aligned}
H^0_d &= \frac{h_d + iP_d}{\sqrt{2}} + \upsilon_d\,, \qquad & H^0_u &= \frac{h_u + iP_u}{\sqrt{2}} + \upsilon_u\,, \\
\tilde{\nu}_i &= \frac{(\tilde{\nu}_i)^R + i(\tilde{\nu}_i)^I}{\sqrt{2}} + \upsilon_{\nu_i}\,, \qquad & \tilde{\nu}^c_i &= \frac{(\tilde{\nu}^c_i)^R + i(\tilde{\nu}^c_i)^I}{\sqrt{2}} + \upsilon_{\nu^c_i}\,. \qquad (5)
\end{aligned}$$

For simplicity we will assume that all parameters in the potential are real in the following. After EWSB, the scalars mass matrices M^2_S, M^2_P and $M^2_{S^\pm}$ are given in Appendix B. The CP-odd neutral scalars mass matrix M^2_P contains a massless unphysical Goldstone boson G^0, which can be written as

$$G^0 = \frac{1}{\sqrt{\upsilon^2_d + \upsilon^2_u + \upsilon_{\nu_i}\upsilon_{\nu_i}}}\Big(\upsilon_dP_d - \upsilon_uP_u + \upsilon_{\nu_i}(\tilde{\nu}_i)^I\Big) \qquad (6)$$

with an 8×8 unitary matrix Z_H

$$Z_H = \begin{pmatrix} \frac{\upsilon_d}{\upsilon_{\text{EW}}} & \frac{\upsilon_u}{\upsilon_{\text{SM}}} & \frac{\upsilon_{\nu_1}\upsilon_d}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & \frac{\upsilon_{\nu_2}\upsilon_d}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & \frac{\upsilon_{\nu_3}\upsilon_d}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & 0_{3\times1} \\ -\frac{\upsilon_u}{\upsilon_{\text{EW}}} & \frac{\upsilon_d}{\upsilon_{\text{SM}}} & -\frac{\upsilon_{\nu_1}\upsilon_u}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & -\frac{\upsilon_{\nu_2}\upsilon_u}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & -\frac{\upsilon_{\nu_3}\upsilon_u}{\upsilon_{\text{EW}}\upsilon_{\text{SM}}} & 0_{3\times1} \\ \frac{\upsilon_{\nu_1}}{\upsilon_{\text{EW}}} & 0 & -\frac{\upsilon_{\text{SM}}}{\upsilon_{\text{EW}}} & \frac{\upsilon_{\nu_3}}{\upsilon_{\text{EW}}} & -\frac{\upsilon_{\nu_2}}{\upsilon_{\text{EW}}} & 0_{3\times1} \\ \frac{\upsilon_{\nu_2}}{\upsilon_{\text{EW}}} & 0 & -\frac{\upsilon_{\nu_3}}{\upsilon_{\text{EW}}} & -\frac{\upsilon_{\text{SM}}}{\upsilon_{\text{EW}}} & \frac{\upsilon_{\nu_1}}{\upsilon_{\text{EW}}} & 0_{3\times1} \\ \frac{\upsilon_{\nu_3}}{\upsilon_{\text{EW}}} & 0 & \frac{\upsilon_{\nu_2}}{\upsilon_{\text{EW}}} & -\frac{\upsilon_{\nu_1}}{\upsilon_{\text{EW}}} & -\frac{\upsilon_{\text{SM}}}{\upsilon_{\text{EW}}} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & 0_{1\times3} & 0_{1\times3} & 0_{1\times3} & 1_{3\times3} \end{pmatrix}, \qquad (7)$$

where $\upsilon_{\text{SM}} = \sqrt{\upsilon_d^2 + \upsilon_u^2}$ and $\upsilon_{\text{EW}} = \sqrt{\upsilon_d^2 + \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}}$. Making use of the minimization conditions of the tree-level neutral scalar potential, which are given in Appendix A, we have

$$\begin{cases} (Z_H^T M_P^2 Z_H)_{11} = 0\,, \\ (Z_H^T M_P^2 Z_H)_{1\alpha} = (Z_H^T M_P^2 Z_H)_{\alpha 1} = 0, \quad \alpha = 2, \ldots, 8. \end{cases} \qquad (8)$$

The remaining 7×7 matrix $\Big((Z_H^T M_P^2 Z_H)_{\alpha\beta}\Big)$ $(\alpha, \beta = 2, \ldots, 8)$ can be further diagonalized, and then gives seven diagonal masses. The charged scalars mass matrix $M_{S^\pm}^2$ also contains the massless unphysical Goldstone bosons $G^\pm$, which can be written as

$$G^\pm = \frac{1}{\sqrt{\upsilon_d^2 + \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}}} \Big(\upsilon_d H_d^\pm - \upsilon_u H_u^\pm + \upsilon_{\nu_i}\tilde{e}_{L_i}^\pm\Big) \qquad (9)$$

with the unitary matrix Z_H and

$$\begin{cases} (Z_H^T M_{S^\pm}^2 Z_H)_{11} = 0\,, \\ (Z_H^T M_{S^\pm}^2 Z_H)_{1\alpha} = (Z_H^T M_{S^\pm}^2 Z_H)_{\alpha 1} = 0, \quad \alpha = 2, \ldots, 8. \end{cases} \qquad (10)$$

In the physical (unitary) gauge, the Goldstone bosons G^0 and $G^\pm$ are eaten by Z-boson and W-boson, respectively, and disappear from the Lagrangian.

Then the mass squared of charged and neutral gauge boson are

$$\begin{cases} m_W^2 = \frac{e^2}{2s_W^2}\Big(\upsilon_u^2 + \upsilon_d^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}\Big), \\ m_Z^2 = \frac{e^2}{2s_W^2 c_W^2}\Big(\upsilon_u^2 + \upsilon_d^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}\Big), \end{cases} \qquad (11)$$

and

$$\tan\beta = \frac{\upsilon_u}{\sqrt{\upsilon_d^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}}} \,. \tag{12}$$

Here e is the electromagnetic coupling constant, $s_{_W} = \sin\theta_{_W}$ and $c_{_W} = \cos\theta_{_W}$ with $\theta_{_W}$ is the Weinberg angle, respectively.

3. Lepton-flavor violation in the $\mu\nu$SSM

In this section, we present the analysis on the decay width of the rare LFV processes $l_j^- \to l_i^-\gamma$ and $l_j^- \to l_i^- l_i^- l_i^+$ in the $\mu\nu$SSM. For this study we will use the indices $\beta, \zeta = 1, \ldots, 5$, $\alpha, \rho = 1, \ldots, 8$, and $\eta, \sigma = 1, \ldots, 10$. And the summation convention is implied on the repeated indices.

3.1. Rare decay $l_j^- \to l_i^-\gamma$



Figure 1: Feynman diagrams for the LFV process $l_j^- \to l_i^-\gamma$. (a) represents the contributions from neutral fermions χ^0_η and charged scalars $S^-_{\alpha,\rho}$ loops, while (b) represents the contributions from charged fermions $\chi_{\beta,\zeta}$ and neutral scalars N_α $(N = S, P)$ loops.

The amplitude for $l_j^- \to l_i^-\gamma$ (including $\mu \to e\gamma$ and $\tau \to \mu\gamma$) is generally written as [18]

$$\begin{aligned} T = e\epsilon^\mu \bar{u}_i(p+q)\Big[&q^2\gamma_\mu(A_1^L P_L + A_1^R P_R) \\ &+ m_{l_j} i\sigma_{\mu\nu}q^\nu(A_2^L P_L + A_2^R P_R)\Big]u_j(p) \,, \end{aligned} \tag{13}$$

where q is the injecting photon momentum, p is the injecting lepton momentum, and m_{l_j} is the mass of the j-th generation charged lepton, respectively. Furthermore, ϵ is the photon polarization vector, $u_i(p)$ ($v_i(p)$ in the expressions below) is the wave function for lepton (antilepton), and $P_L = \frac{1}{2}(1-\gamma_5)$,

$P_R = \frac{1}{2}(1+\gamma_5)$. Here, the Feynman diagrams contributing to the above amplitude are shown in Fig.1. And the coefficients can be written by

$$A_a^{L,R} = A_a^{(n)L,R} + A_a^{(c)L,R} \quad (a=1,2)\;, \tag{14}$$

where $A_a^{(n)L,R}$ denote the contributions from the virtual neutral fermion loops, and $A_a^{(c)L,R}$ stand for the contributions from the virtual charged fermion loops, respectively. After integrating the heavy freedoms out, we formulate those coefficients as follows

$$\begin{aligned}
A_1^{(n)L} &= \frac{1}{6m_W^2} C_R^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_i} C_L^{S_\alpha^{-*} \chi_j \bar{\chi}_\eta^\circ} I_4(x_{\chi_\eta^\circ}, x_{S_\alpha^-})\;,\\
A_2^{(n)L} &= \frac{m_{\chi_\eta^\circ}}{m_{l_j} m_W^2} C_L^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_i} C_L^{S_\alpha^{-*} \chi_j \bar{\chi}_\eta^\circ} \Big[I_3(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) - I_1(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) \Big]\;,\\
A_a^{(n)R} &= A_a^{(n)L}\Big|_{L\leftrightarrow R}\;,
\end{aligned} \tag{15}$$

where the concrete expressions for form factors I_k $(k=1,\ldots,4)$ can be found in Appendix E. Additionally, $x = m^2/m_W^2$, m is the mass for the corresponding particle and m_W is the mass for the W-boson, respectively. In a similar way, the corrections from the Feynman diagrams with virtual charged fermions are

$$\begin{aligned}
A_1^{(c)L} &= \sum_{N=S,P} \frac{1}{6m_W^2} C_R^{N_\alpha \chi_\beta \bar{\chi}_i} C_L^{N_\alpha \chi_j \bar{\chi}_\beta} \Big[I_1(x_{\chi_\beta}, x_{N_\alpha}) - 2I_2(x_{\chi_\beta}, x_{N_\alpha}) \\
&\qquad - I_4(x_{\chi_\beta}, x_{N_\alpha}) \Big]\;,\\
A_2^{(c)L} &= \sum_{N=S,P} \frac{m_{\chi_\beta}}{m_{l_j} m_W^2} C_L^{N_\alpha \chi_\beta \bar{\chi}_i} C_L^{N_\alpha \chi_j \bar{\chi}_\beta} \Big[I_1(x_{\chi_\beta}, x_{N_\alpha}) - I_2(x_{\chi_\beta}, x_{N_\alpha}) \\
&\qquad - I_4(x_{\chi_\beta}, x_{N_\alpha}) \Big]\;,\\
A_a^{(c)R} &= A_a^{(c)L}\Big|_{L\leftrightarrow R}\;.
\end{aligned} \tag{16}$$

Using the amplitude presented in Eq.(13), we then obtain the decay width for $l_j^- \to l_i^- \gamma$ as [18]

$$\Gamma(l_j^- \to l_i^- \gamma) = \frac{e^2}{16\pi} m_{l_j}^5 \left(\left|A_2^L\right|^2 + \left|A_2^R\right|^2 \right)\;. \tag{17}$$

And the branching ratio of $l_j^- \to l_i^- \gamma$ is

$$\mathrm{Br}(l_j^- \to l_i^- \gamma) = \frac{\Gamma(l_j^- \to l_i^- \gamma)}{\Gamma_{l_j^-}}\;, \tag{18}$$

where $\Gamma_{l_j^-}$ denotes the total decay rate of the lepton l_j^-. In the numerical calculation, $\Gamma_\mu \approx 2.996\times10^{-19}\,\mathrm{GeV}$ for the muon and $\Gamma_\tau \approx 2.265\times10^{-12}\,\mathrm{GeV}$ for the tauon.

3.2. Rare decay $l_j^- \to l_i^- l_i^- l_i^+$



Figure 2: Penguin-type diagrams for the LFV process $l_j^- \to l_i^- l_i^- l_i^+$ in which a photon γ and Z-boson are exchanged. The blob indicates an $l_j^- - l_i^- - \gamma$ vertex such as Fig.1 or $l_j^- - l_i^- - Z$ vertex where the Z-boson is external.

For the rare LFV processes $l_j^- \to l_i^- l_i^- l_i^+$ (including $\mu \to 3e$), the corresponding effective Hamilton originates from penguin-type diagrams and from box-type diagrams. The γ-penguin contribution can be computed using Eq.(13), with the result

$$\begin{aligned} T_{\gamma-\mathrm{p}} = {}& \bar{u}_i(p_1)\Big[q^2\gamma_\mu(A_1^L P_L + A_1^R P_R) + m_{l_j} i\sigma_{\mu\nu}q^\nu(A_2^L P_L + A_2^R P_R)\Big]u_j(p) \\ & \times \frac{e^2}{q^2}\bar{u}_i(p_2)\gamma^\mu v_i(p_3) - (p_1 \leftrightarrow p_2)\,. \end{aligned} \tag{19}$$

Similarly, the contribution from Z-penguin diagrams which are depicted by Fig.2 is

$$\begin{aligned} T_{Z-\mathrm{p}} = {}& \frac{e^2}{m_Z^2}\bar{u}_i(p_1)\gamma_\mu(F_L P_L + F_R P_R)u_j(p)\bar{u}_i(p_2)\gamma^\mu\Big(C_L^{Z\chi_{2+i}\bar{\chi}_{2+i}}P_L \\ & + C_R^{Z\chi_{2+i}\bar{\chi}_{2+i}}P_R\Big)v_i(p_3) - (p_1 \leftrightarrow p_2)\,, \end{aligned} \tag{20}$$

where m_Z is the mass for the Z-boson and

$$F_{L,R} = F_{L,R}^{(n)} + F_{L,R}^{(c)}\,. \tag{21}$$

The contributions to the effective couplings $F_{L,R}^{(n)}$ and $F_{L,R}^{(c)}$ are

$$\begin{aligned}
F_L^{(n)} =& \sum_{N=S,P} \Big[\frac{m_{\chi_\zeta} m_{\chi_\beta}}{e^2 m_W^2} C_R^{N_\alpha \chi_\zeta \bar{\chi}_i} C_L^{Z \chi_\beta \bar{\chi}_\zeta} C_L^{N_\alpha \chi_j \bar{\chi}_\beta} G_1(x_{N_\alpha}, x_{\chi_\zeta}, x_{\chi_\beta}) \\
& - \frac{1}{2e^2} C_R^{N_\alpha \chi_\zeta \bar{\chi}_i} C_R^{Z \chi_\beta \bar{\chi}_\zeta} C_L^{N_\alpha \chi_j \bar{\chi}_\beta} G_2(x_{N_\alpha}, x_{\chi_\zeta}, x_{\chi_\beta})\Big] , \\
F_L^{(c)} =& \frac{1}{2e^2} C_R^{S_\rho^- \chi_\eta^0 \bar{\chi}_i} C_R^{Z S_\alpha^- S_\rho^{-*}} C_L^{S_\alpha^{-*} \chi_j \bar{\chi}_\eta^0} G_2(x_{\chi_\eta^0}, x_{S_\alpha^-}, x_{S_\rho^-}) , \\
F_R^{(n,c)} =& F_L^{(n,c)}\Big|_{L \leftrightarrow R} . \qquad (22)
\end{aligned}$$

Here, the concrete expressions for G_k are given in Appendix E.



Figure 3: Box-type diagrams for the LFV process $l_j^- \to l_i^- l_i^- l_i^+$. (a) represents the contributions from neutral fermions $\chi_{\eta,\sigma}^0$ and charged scalars $S_{\alpha,\rho}^-$ loops, and (b) represents the contributions from charged fermions $\chi_{\beta,\zeta}$ and neutral scalars $N_{\alpha,\rho}$ $(N = S, P)$ loops.

Furthermore, the effective Hamilton from the box-type diagrams which are drawn in Fig.3 can be written as

$$\begin{aligned}
T_{box} =& \Big\{B_1^L e^2 \bar{u}_i(p_1)\gamma_\mu P_L u_j(p) \bar{u}_i(p_2)\gamma^\mu P_L v_i(p_3) + (L \leftrightarrow R)\Big\} \\
& + \Big\{B_2^L e^2 \Big[\bar{u}_i(p_1)\gamma_\mu P_L u_j(p) \bar{u}_i(p_2)\gamma^\mu P_R v_i(p_3) - (p_1 \leftrightarrow p_2)\Big] + (L \leftrightarrow R)\Big\} \\
& + \Big\{B_3^L e^2 \Big[\bar{u}_i(p_1) P_L u_j(p) \bar{u}_i(p_2) P_L v_i(p_3) - (p_1 \leftrightarrow p_2)\Big] + (L \leftrightarrow R)\Big\}
\end{aligned}$$

$$+\Big\{B_4^L e^2\Big[\bar{u}_i(p_1)\sigma_{\mu\nu}P_L u_j(p)\bar{u}_i(p_2)\sigma^{\mu\nu}P_L v_i(p_3)-(p_1\leftrightarrow p_2)\Big]+\;(L\leftrightarrow R)\Big\} \tag{23}$$

with

$$B_a^{L,R}=B_a^{(n)L,R}+B_a^{(c)L,R}\quad(a=1,\ldots,4)\,. \tag{24}$$

The effective couplings $B_a^{(n)L,R}$ originate from those box diagrams with virtual neutral fermion contributions:

$$\begin{aligned}
B_1^{(n)L}=&\;\frac{m_{\chi_\eta^0}m_{\chi_\sigma^0}}{e^2m_W^4}G_3(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})C_L^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_R^{S_\rho^-\chi_\sigma^0\bar{\chi}_i}C_R^{S_\alpha^{-*}\chi_i\bar{\chi}_\sigma^0}\\
&+\frac{1}{2e^2m_W^2}G_4(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})\Big[C_R^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_R^{S_\alpha^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\rho^{-*}\chi_i\bar{\chi}_\sigma^0}\\
&+C_L^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_R^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_R^{S_\rho^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_i\bar{\chi}_\sigma^0}\Big],\\
B_2^{(n)L}=&-\frac{m_{\chi_\eta^0}m_{\chi_\sigma^0}}{2e^2m_W^4}G_3(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})C_R^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_R^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_L^{S_\alpha^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\rho^{-*}\chi_i\bar{\chi}_\sigma^0}\\
&+\frac{1}{4e^2m_W^2}G_4(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})\Big[C_R^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_L^{S_\alpha^-\chi_\sigma^0\bar{\chi}_i}C_R^{S_\rho^{-*}\chi_i\bar{\chi}_\sigma^0}\\
&+C_R^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_R^{S_\rho^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_i\bar{\chi}_\sigma^0}\Big],\\
B_3^{(n)L}=&\;\frac{m_{\chi_\eta^0}m_{\chi_\sigma^0}}{e^2m_W^4}G_3(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})\Big[C_L^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_L^{S_\alpha^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\rho^{-*}\chi_i\bar{\chi}_\sigma^0}\\
&-\frac{1}{2}\,C_L^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_L^{S_\rho^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_i\bar{\chi}_\sigma^0}\Big],\\
B_4^{(n)L}=&\;\frac{m_{\chi_\eta^0}m_{\chi_\sigma^0}}{8e^2m_W^4}G_3(x_{\chi_\eta^0},x_{\chi_\sigma^0},x_{S_\alpha^-},x_{S_\rho^-})C_L^{S_\rho^-\chi_\eta^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_j\bar{\chi}_\eta^0}C_L^{S_\rho^-\chi_\sigma^0\bar{\chi}_i}C_L^{S_\alpha^{-*}\chi_i\bar{\chi}_\sigma^0},\\
B_a^{(n)R}=&\;B_a^{(n)L}\Big|_{L\leftrightarrow R}\,.
\end{aligned} \tag{25}$$

Correspondingly, the effective couplings from the box diagrams with virtual charged fermion contributions $B_a^{(c)L,R}$ are

$$\begin{aligned}
B_1^{(c)L}=&\sum_{N=S,P}\frac{1}{2e^2m_W^2}G_4(x_{\chi_\zeta},x_{\chi_\beta},x_{N_\alpha},x_{N_\rho})C_R^{N_\rho\chi_\zeta\bar{\chi}_i}C_L^{N_\alpha\chi_j\bar{\chi}_\zeta}C_R^{N_\alpha\chi_\beta\bar{\chi}_i}C_L^{N_\rho\chi_i\bar{\chi}_\beta},\\
B_2^{(c)L}=&\sum_{N=S,P}\Big[\frac{1}{4e^2m_W^2}G_4(x_{\chi_\zeta},x_{\chi_\beta},x_{N_\alpha},x_{N_\rho})C_R^{N_\rho\chi_\zeta\bar{\chi}_i}C_L^{N_\alpha\chi_j\bar{\chi}_\zeta}C_L^{N_\alpha\chi_\beta\bar{\chi}_i}C_R^{N_\rho\chi_i\bar{\chi}_\beta}\\
&-\frac{m_{\chi_\zeta}m_{\chi_\beta}}{2e^2m_W^4}G_3(x_{\chi_\zeta},x_{\chi_\beta},x_{N_\alpha},x_{N_\rho})C_R^{N_\rho\chi_\zeta\bar{\chi}_i}C_R^{N_\alpha\chi_j\bar{\chi}_\zeta}C_L^{N_\alpha\chi_\beta\bar{\chi}_i}C_L^{N_\rho\chi_i\bar{\chi}_\beta}\Big],
\end{aligned}$$

$$
\begin{aligned}
B_3^{(c)L} &= \sum_{N=S,P} \frac{m_{\chi_\zeta} m_{\chi_\beta}}{e^2 m_W^4} G_3(x_{\chi_\zeta}, x_{\chi_\beta}, x_{N_\alpha}, x_{N_\rho}) C_L^{N_\rho \chi_\zeta \bar{\chi}_i} C_L^{N_\alpha \chi_j \bar{\chi}_\zeta} C_L^{N_\alpha \chi_\beta \bar{\chi}_i} C_L^{N_\rho \chi_i \bar{\chi}_\beta} , \\
B_4^{(c)L} &= 0 , \\
B_a^{(c)R} &= B_a^{(c)L}\Big|_{L\leftrightarrow R} .
\end{aligned}
\tag{26}
$$

Using the expression for the above amplitude, we can calculate the decay width for $l_j^- \to l_i^- l_i^- l_i^+$ [18]:

$$
\begin{aligned}
\Gamma(l_j^- \to l_i^- l_i^- l_i^+) &= \frac{e^4}{512\pi^3} m_{l_j}^5 \Big\{ (\left|A_2^L\right|^2 + \left|A_2^R\right|^2)(\frac{16}{3}\ln\frac{m_{l_j}}{2m_{l_i}} - \frac{14}{9}) \\
&+ (\left|A_1^L\right|^2 + \left|A_1^R\right|^2) - 2(A_1^L A_2^{R*} + A_2^L A_1^{R*} + \text{H.c.}) + \frac{1}{6}(\left|B_1^L\right|^2 + \left|B_1^R\right|^2) \\
&+ \frac{1}{3}(\left|B_2^L\right|^2 + \left|B_2^R\right|^2) + \frac{1}{24}(\left|B_3^L\right|^2 + \left|B_3^R\right|^2) + 6(\left|B_4^L\right|^2 + \left|B_4^R\right|^2) \\
&- \frac{1}{2}(B_3^L B_4^{L*} + B_3^R A_4^{R*} + \text{H.c.}) + \frac{1}{3}(A_1^L B_1^{L*} + A_1^R B_1^{R*} + A_1^L B_2^{L*} \\
&+ A_1^R B_2^{R*} + \text{H.c.}) - \frac{2}{3}(A_2^R B_1^{L*} + A_2^L B_1^{R*} + A_2^L B_2^{R*} + A_2^R B_2^{L*} + \text{H.c.}) \\
&+ \frac{1}{3}\Big[2(|F_{LL}|^2 + |F_{RR}|^2) + (|F_{LR}|^2 + |F_{RL}|^2) + (B_1^L F_{LL}^* + B_1^R F_{RR}^* \\
&+ B_2^L F_{LR}^* + B_2^R F_{RL}^* + \text{H.c.}) + 2(A_1^L F_{LL}^* + A_1^R F_{RR}^* + \text{H.c.}) \\
&+ (A_1^L F_{LR}^* + A_1^R F_{RL}^* + \text{H.c.}) - 4(A_2^R F_{LL}^* + A_2^L F_{RR}^* + \text{H.c.}) \\
&- 2(A_2^L F_{RL}^* + A_2^R F_{LR}^* + \text{H.c.})\Big]\Big\}
\end{aligned}
\tag{27}
$$

with

$$
\begin{aligned}
F_{LL} &= \frac{F_L C_L^{Z\chi_{2+i}\bar{\chi}_{2+i}}}{m_Z^2}, \qquad F_{RR} = F_{LL}\,|_{L\leftrightarrow R} , \\
F_{LR} &= \frac{F_L C_R^{Z\chi_{2+i}\bar{\chi}_{2+i}}}{m_Z^2}, \qquad F_{RL} = F_{LR}\,|_{L\leftrightarrow R} .
\end{aligned}
\tag{28}
$$

And the branching ratio of $l_j^- \to l_i^- l_i^- l_i^+$ is

$$
\text{Br}(l_j^- \to l_i^- l_i^- l_i^+) = \frac{\Gamma(l_j^- \to l_i^- l_i^- l_i^+)}{\Gamma_{l_j^-}} . \tag{29}
$$

4. $(g-2)_\mu$ in the $\mu\nu$SSM

The anomalous magnetic dipole moment (MDM) of the muon can be actually be written as the operator

$$\mathcal{L}_{MDM} = \frac{e}{4m_\mu} a_\mu \bar{l}_\mu \sigma^{\alpha\beta} l_\mu F_{\alpha\beta}\;, \tag{30}$$

where $\sigma^{\alpha\beta} = \frac{i}{2}[\gamma^\alpha, \gamma^\beta]$, $F_{\alpha\beta}$ is the electromagnetic field strength, l_μ denotes the muon which is on-shell, m_μ is the muon mass and $a_\mu = \frac{1}{2}(g-2)_\mu$. Adopting the effective Lagrangian approach, we can get [19]

$$a_\mu = \frac{4Q_f m_\mu^2}{(4\pi)^2} \Re(C_2^R + C_2^{L*} + C_6^R)\;, \tag{31}$$

where $Q_f = -1$, $\Re(\cdots)$ represents the operation to take the real part of a complex number and $C_{2,6}^{L,R}$ denote the Wilson coefficients of the corresponding operators $O_{2,6}^{L,R}$

$$\begin{aligned}
O_2^{L,R} &= \frac{eQ_f}{(4\pi)^2} \overline{(i\mathcal{D}_\alpha l_\mu)} \gamma^\alpha F \cdot \sigma P_{L,R} l_\mu\;, \\
O_6^{L,R} &= \frac{eQ_f m_\mu}{(4\pi)^2} \bar{l}_\mu F \cdot \sigma P_{L,R} l_\mu\;.
\end{aligned} \tag{32}$$

In the $\mu\nu$SSM, the SUSY corrections can be written as

$$C_{2,6}^{L,R} = C_{2,6}^{L,R(n)} + C_{2,6}^{L,R(c)}\;. \tag{33}$$

The effective couplings $C_{2,6}^{L,R(n)}$ represent the contributions from the triangle diagrams with virtual neutralinos

$$\begin{aligned}
C_2^{R(n)} &= \frac{(4\pi)^2}{Q_f m_W^2} C_L^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_4} C_R^{S_\alpha^{-*} \chi_4 \bar{\chi}_\eta^\circ} \Big[- I_3(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) + I_4(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) \Big]\;, \\
C_6^{R(n)} &= \frac{(4\pi)^2 m_{\chi_\eta^\circ}}{Q_f m_W^2 m_\mu} C_R^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_4} C_R^{S_\alpha^{-*} \chi_4 \bar{\chi}_\eta^\circ} \Big[- 2I_1(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) + 2I_3(x_{\chi_\eta^\circ}, x_{S_\alpha^-}) \Big]\;, \\
C_{2,6}^{L(n)} &= C_{2,6}^{R(n)} \;|_{L\leftrightarrow R}\;\;.
\end{aligned} \tag{34}$$

Similarly, the contributions $C_{2,6}^{L,R(c)}$ originating from triangle diagrams with virtual charginos are

$$\begin{aligned}
C_2^{R(c)} &= \sum_{N=S,P} \frac{(4\pi)^2}{Q_f m_W^2} C_R^{N_\alpha \chi_\beta \bar{\chi}_4} C_L^{N_\alpha \chi_4 \bar{\chi}_\beta} \Big[- I_1(x_{\chi_\beta}, x_{N_\alpha}) + 2I_3(x_{\chi_\beta}, x_{N_\alpha}) \\
&\qquad - I_4(x_{\chi_\beta}, x_{N_\alpha}) \Big] , \\
C_6^{R(c)} &= \sum_{N=S,P} \frac{(4\pi)^2 m_{\chi_\beta}}{Q_f m_W^2 m_\mu} C_R^{N_\alpha \chi_\beta \bar{\chi}_4} C_R^{N_\alpha \chi_4 \bar{\chi}_\beta} \Big[2I_1(x_{\chi_\beta}, x_{N_\alpha}) - 2I_2(x_{\chi_\beta}, x_{N_\alpha}) \\
&\qquad - 2I_3(x_{\chi_\beta}, x_{N_\alpha}) \Big] , \\
C_{2,6}^{L(c)} &= C_{2,6}^{R(c)} \, |_{L \leftrightarrow R} \ . \qquad (35)
\end{aligned}$$

5. The numerical results

5.1. The parameter space

It is well known that there are many free parameters in various SUSY extensions of the SM. In order to obtain a more transparent numerical results, we take some assumptions on parameter space of the $\mu\nu$SSM before we perform the numerical analysis.

In lepton sector, we adopt the minimal flavor violation (MFV) assumptions

$$\begin{aligned}
&\kappa_{ijk} = \kappa \text{ and } (A_\kappa \kappa)_{ijk} = A_\kappa \kappa, \text{ if } i = j = k, \text{ and zero otherwise,} \\
&m^2_{\tilde{L}_{ij}} = m^2_{\tilde{L}_i} \delta_{ij},\ m^2_{\tilde{\nu}^c_{ij}} = m^2_{\tilde{\nu}^c_i} \delta_{ij},\ m^2_{\tilde{e}^c_{ij}} = m^2_{\tilde{e}^c} \delta_{ij}, \\
&Y_{\nu_{ij}} = Y_{\nu_i} \delta_{ij},\ Y_{e_{ij}} = Y_{e_i} \delta_{ij},\ \lambda_i = \lambda,\ \upsilon_{\nu^c_i} = \upsilon_{\nu^c}, \\
&(A_\nu Y_\nu)_{ij} = A_\nu Y_{\nu_i} \delta_{ij},\ (A_e Y_e)_{ij} = A_e Y_{e_i} \delta_{ij} \text{ , and } (A_\lambda \lambda)_i = A_\lambda \lambda, \qquad (36)
\end{aligned}$$

where $i,\ j,\ k = 1,\ 2,\ 3$.

The 3×3 matrix Y_ν determines the Dirac masses for the neutrinos $Y_\nu \upsilon_u \sim m_D$, and the tiny neutrino masses are obtained through TeV scale seesaw mechanism $m_\nu \sim m_D m_N^{-1} m_D^T$. This indicates that the nonzero VEVs of left-handed sneutrinos satisfy $\upsilon_{\nu_i} \ll \upsilon_{u,d}$, then

$$\tan\beta \simeq \frac{\upsilon_u}{\upsilon_d} \ . \qquad (37)$$

Assuming that the charged lepton mass matrix in the flavor basic is in the diagonal form, we get

$$Y_{e_i} = \frac{m_{l_i}}{\upsilon_d}, \tag{38}$$

where m_{l_i} is the charged lepton l_i mass, and we parameterize the unitary matrix which diagonalizes the effective light neutrino mass matrix m_{eff} (can be found in Appendix C) as [20]

$$\begin{aligned} U_\nu = & \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \\ & \times diag(1, e^{i\frac{\alpha_{21}}{2}}, e^{i\frac{\alpha_{31}}{2}}) \,, \end{aligned} \tag{39}$$

where $c_{ij} = \cos\theta_{ij}$, $s_{ij} = \sin\theta_{ij}$, the angles $\theta_{ij} = [\,0, \pi/2\,]$, $\delta = [\,0, 2\pi\,]$ is the Dirac CP violation phase and α_{21}, α_{31} are two Majorana CP violation phases. Here, we choose $\delta = \alpha_{21} = \alpha_{31} = 0$. U_ν diagonalizes m_{eff} in the following way:

$$U_\nu^T m_{eff}^T m_{eff} U_\nu = diag(m_{\nu_1}^2, m_{\nu_2}^2, m_{\nu_3}^2) \,, \tag{40}$$

where the neutrino mass m_{ν_i} connected with experimental measurements through

$$m_{\nu_2}^2 - m_{\nu_1}^2 = \Delta m_{21}^2, \qquad m_{\nu_3}^2 - m_{\nu_2}^2 = \Delta m_{32}^2. \tag{41}$$

The combination of Eq.(39), Eq.(40), Eq.(41) with neutrino oscillation experimental data gives some strong constraints on relevant parameter space of the $\mu\nu$SSM.

At the EW scale, the soft masses $m_{\tilde{H}_d}^2$, $m_{\tilde{H}_u}^2$, $m_{\tilde{L}_i}^2$ and $m_{\tilde{\nu}_i^c}^2$ are derived from the minimization conditions of the tree-level neutral scalar potential, which are given in Appendix A. Implying the approximate GUT relation $M_1 = \frac{\alpha_1^2}{\alpha_2^2} M_2 \approx 0.5 M_2$, the free parameters affect our analysis are

$$\lambda,\ \kappa,\ \tan\beta,\ A_{\lambda,\kappa,\nu,e},\ m_{\tilde{e}^c},\ \upsilon_{\nu^c},\ M_2\ . \tag{42}$$

To obtain the Yukawa couplings Y_{ν_i} and υ_{ν_i} from Eq.(40), we assume the neutrinos masses satisfying $m_{\nu_1} < m_{\nu_2} < m_{\nu_3}$, and choose $m_{\nu_2} = 10^{-2}\,\text{eV}$ as input in our numerical analysis. Then we can get $m_{\nu_{1,3}}$ from the experimental data on the differences of neutrino mass squared. For U_ν, the values of θ_{ij} are obtained from the experimental data in Eq.(1). And the effective light neutrino mass matrix m_{eff} can approximate as [16]

$$m_{eff_{ij}} \approx \frac{2A\upsilon_{\nu^c}}{3\Delta} b_i b_j + \frac{1-3\delta_{ij}}{6\kappa\upsilon_{\nu^c}} a_i a_j \; , \tag{43}$$

where

$$\begin{aligned}
&\Delta = \lambda^2 (\upsilon_d^2 + \upsilon_u^2)^2 + 4\lambda\kappa\upsilon_{\nu^c}^2 \upsilon_d \upsilon_u - 12\lambda^2 \upsilon_{\nu^c} AB \; , \\
&A = \kappa\upsilon_{\nu^c}^2 + \lambda\upsilon_d\upsilon_u \; , \\
&\frac{1}{B} = \frac{e^2}{c_W^2 M_1} + \frac{e^2}{s_W^2 M_2} \; , \\
&a_i = Y_{\nu_i}\upsilon_u \; , \quad b_i = Y_{\nu_i}\upsilon_d + 3\lambda\upsilon_{\nu_i} \; .
\end{aligned} \tag{44}$$

Then, we can numerically derive $Y_{\nu_i} \sim \mathcal{O}(10^{-7})$ and $\upsilon_{\nu_i} \sim \mathcal{O}(10^{-4}\text{GeV})$ from Eq.(40).

5.2. Branching ratio of LFV processes



Figure 4: Branching ratio for the process $\mu \to e\gamma$ varies with M_2 for $\tan\beta = 3, \; 10, \; 30$, respectively.

Considering the research of the $\mu\nu$SSM [4], we choose the relevant parameters as $\lambda = 0.1$, $\kappa = 0.01$, $m_{\tilde{e}^c} = A_e = A_\lambda = 1$ TeV, $A_\nu = A_\kappa = -1$ TeV and $\upsilon_{\nu^c} = 800$ GeV in next numerical analysis for convenience. With those assumptions on parameter space, we present the branching ratio of $\mu \to e\gamma$



Figure 5: Branching ratio for the process $\mu \to 3e$ varies with M_2 for $\tan\beta = 3,\ 10,\ 30$, respectively.

versus M_2 in Fig.4. As $M_2 \leq 2$ TeV, the theoretical evaluations exceed the upper experimental bound easily. The fact implies that experimental data do not favor small M_2. Along with increasing of M_2, theoretical evaluation on the branching ratio of $\mu \to e\gamma$ decreases steeply. As $M_2 = 3$ TeV and $\tan\beta = 10$, theoretical evaluation on the branching ratio of $\mu \to e\gamma$ is about 5×10^{-13} which can be detected in near future. In the future, the expected sensitivity for Br($\mu \to e\gamma$) would be of order 10^{-13} [21]. Differing from LFV processes which are researched in the BRpV model [22], the large VEVs of right-handed sneutrinos in the $\mu\nu$SSM induce new sources for lepton-flavor violation. So, here the branching ratio of $\mu \to e\gamma$ can easily reach the upper experimental bound 2.4×10^{-12} [13].

We also investigate the $\mu \to 3e$ processes in detail. And the branching ratio of $\mu \to 3e$ is also decreases with increasing of M_2, and raises with increasing of $\tan\beta$, which is presented in the Fig.5. By Introducing the right-handed sneutrinos which the VEVs are nonzero to the $\mu\nu$SSM, the branching ratio of $\mu \to 3e$ can also easily reach the upper experimental bound 10^{-12} [13]. We can see that the experimental bounds of the branching ratio of $\mu \to 3e$ and $\mu \to e\gamma$ give very strong constraints on the $\mu\nu$SSM.

In Fig.6, we show the branching ratio for $\tau \to \mu\gamma$ versus M_2 as $\tan\beta = 3,\ 10,\ 30$. Similar to the case of $\mu \to e\gamma$, the evaluation on the branching ratio for $\tau \to \mu\gamma$ decreases with increasing of M_2, and is enhanced by large $\tan\beta$. As $M_2 = 3$ TeV and $\tan\beta = 10$, Br($\tau \to \mu\gamma$) $\approx 10^{-13}$ is four orders below the expected sensitivity 10^{-9} [23].



Figure 6: Branching ratio for the process $\tau \to \mu\gamma$ varies with M_2 for $\tan\beta = 3,\ 10,\ 30$, respectively.

5.3. Muon anomalous magnetic dipole moment



Figure 7: The SUSY contribution to the anomalous magnetic dipole moment of the muon varies with M_2 for $\tan\beta = 3,\ 10,\ 30$, respectively. The gray area denotes the Δa_μ at 1.8 standard deviation.

Finally, we analyze the anomalous magnetic dipole moment of the muon in the $\mu\nu$SSM. Rescaled the final result of the E821 Collaboration at BNL [24] using μ/p magnetic moment ratio of 3.183345137(85) from ref.[25], the PDG Collaboration [13] gives the world average of muon anomalous magnetic dipole moment

$$a_\mu^{\mathrm{exp}} = \frac{1}{2}(g_\mu - 2) = 11659208.9(5.4)(3.3) \times 10^{-10}\,, \tag{45}$$

where the statistical and systematic uncertainties are given, respectively. And the Standard Model (SM) prediction [13] is

$$a_\mu^{\rm SM} = 11659184.1(4.8) \times 10^{-10}\,. \tag{46}$$

So, the difference between experiment and the SM prediction

$$\Delta a_\mu = a_\mu^{\rm exp} - a_\mu^{\rm SM} = 24.8(8.7)(4.8) \times 10^{-10}\,, \tag{47}$$

represents an interesting but not yet conclusive discrepancy of 1.8 standard deviation. An alternate interpretation is that Δa_μ may be a new physics signal with supersymmetric particle loops as the leading candidate explanation. If treated the supersymmetry as the leading explanation, parameter space of the $\mu\nu$SSM should be constrained by the experimental data on Δa_μ.

The SUSY contribution to the muon anomalous magnetic dipole moment in the $\mu\nu$SSM is shown in Fig.7. The result shows that when $\tan\beta = 3$, Δa_μ constrains $M_2 < 1$ TeV, which is opposite to what the upper experimental bound of ${\rm Br}(\mu \rightarrow {\rm e}\gamma)$ constrains. The fact implies that experimental data do not favor small $\tan\beta$ in the $\mu\nu$SSM with the MFV assumptions (36). When $\tan\beta = 30$, Δa_μ constrains $2\ {\rm TeV} \leq M_2 < 7$ TeV, compared with that the upper experimental bound of ${\rm Br}(\mu \rightarrow {\rm e}\gamma)$ constrains $M_2 \geq 3.5$ TeV, the M_2 has more consistent interval. So, under the MFV assumptions, the $\mu\nu$SSM favors large $\tan\beta$ and M_2 for consistent with experimental data.

6. Conclusions

Besides the superfields of the MSSM, the $\mu\nu$SSM introduces three exotic right-handed sneutrinos $\hat{\nu}_i^c$ to solve the μ problem of the MSSM. And exotic right-handed sneutrinos which the vacuum expectation values are nonzero induce new sources for lepton-flavor violation. In addition, from the scalars for the $\mu\nu$SSM we strictly separate the Goldstone bosons, which disappear in the physical gauge.

Considering the updated experimental data on neutrino oscillations, we analyze various LFV processes and $(g-2)_\mu$ in the $\mu\nu$SSM. Numerical results indicate that the new physics corrections dominate the evaluation on the branching ratios of LFV processes in some parameter space of the $\mu\nu$SSM. And the theoretical predictions on the branching ratios of LFV processes $\mu \rightarrow e\gamma$ and $\mu \rightarrow 3e$ for large $\tan\beta$ can easily reach the present experimental upper bounds and be detected in near future. Additionally, the present experimental observations on $(g-2)_\mu$ also give very strong constraint on the model. Under the MFV assumptions (36), the $\mu\nu$SSM favors large $\tan\beta$ and M_2 for consistent with experimental data. Certainly, a neutral Higgs with mass $m_{h_0} \sim 124 - 126$ GeV reported by ATLAS [26] and CMS [27] also contributes a strict constraint on relevant parameter space, we will discuss this problem elsewhere.

Acknowledgements

The work has been supported by the National Natural Science Foundation of China (NNSFC) with Grant No. 10975027, No. 11275036, No. 11047002 and Natural Science Fund of Hebei University with Grant No. 2011JQ05, No. 2012-242.

Appendix A. Minimization of the potential

First, the eight minimization conditions of the tree-level neutral scalar potential are given below:

$$\begin{aligned}
&m_{H_d}^2 \upsilon_d + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i})\upsilon_d - (A_\lambda\lambda)_i \upsilon_u \upsilon_{\nu_i^c} - \lambda_j \kappa_{ijk} \upsilon_u \upsilon_{\nu_i^c} \upsilon_{\nu_k^c} \\
&\quad + (\lambda_i\lambda_j \upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + \lambda_i\lambda_i \upsilon_u^2)\upsilon_d - Y_{\nu_{ij}} \upsilon_{\nu_i} (\lambda_k \upsilon_{\nu_k^c}\upsilon_{\nu_j^c} + \lambda_j \upsilon_u^2) = 0\,,
\end{aligned} \tag{A.1}$$

$$\begin{aligned}
&m_{H_u}^2 \upsilon_u - \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i})\upsilon_u + (A_\nu Y_\nu)_{ij} \upsilon_{\nu_i}\upsilon_{\nu_j^c} - (A_\lambda\lambda)_i \upsilon_d \upsilon_{\nu_i^c} \\
&\quad + (\lambda_i\lambda_j \upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + \lambda_i\lambda_i \upsilon_u^2)\upsilon_u + Y_{\nu_{ij}} \upsilon_{\nu_i} (\kappa_{ljk} \upsilon_{\nu_l^c}\upsilon_{\nu_k^c} - 2\lambda_j \upsilon_d \upsilon_u) \\
&\quad - \lambda_j \kappa_{ijk} \upsilon_d \upsilon_{\nu_i^c}\upsilon_{\nu_k^c} + (Y_{\nu_{ki}} Y_{\nu_{kj}} \upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + Y_{\nu_{ik}} Y_{\nu_{jk}} \upsilon_{\nu_i}\upsilon_{\nu_j})\upsilon_u = 0\,,
\end{aligned} \tag{A.2}$$

$$\begin{aligned}
&m_{\tilde{L}_{ij}}^2 \upsilon_{\nu_j} + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_j}\upsilon_{\nu_j})\upsilon_{\nu_i} + (A_\nu Y_\nu)_{ij} \upsilon_u \upsilon_{\nu_j^c} + Y_{\nu_{il}} \kappa_{ljk} \upsilon_u \upsilon_{\nu_j^c}\upsilon_{\nu_k^c} \\
&\quad - Y_{\nu_{ij}} \lambda_k \upsilon_{\nu_j^c}\upsilon_{\nu_k^c}\upsilon_d - Y_{\nu_{ij}} \lambda_j \upsilon_u^2 \upsilon_d + Y_{\nu_{ij}} Y_{\nu_{lk}} \upsilon_{\nu_l} \upsilon_{\nu_j^c}\upsilon_{\nu_k^c} \\
&\quad + Y_{\nu_{ik}} Y_{\nu_{jk}} \upsilon_u^2 \upsilon_{\nu_j} = 0\,,
\end{aligned} \tag{A.3}$$

$$\begin{aligned}
&m_{\tilde{\nu}_{ij}^c}^2 \upsilon_{\nu_j^c} + (A_\nu Y_\nu)_{ji} \upsilon_{\nu_j} \upsilon_u - (A_\lambda\lambda)_i \upsilon_d \upsilon_u + (A_\kappa\kappa)_{ijk} \upsilon_{\nu_j^c}\upsilon_{\nu_k^c} - 2\lambda_j \kappa_{ijk} \upsilon_d \upsilon_u \upsilon_{\nu_k^c} \\
&\quad + \lambda_i\lambda_j \upsilon_{\nu_j^c}(\upsilon_d^2 + \upsilon_u^2) + 2\kappa_{lim}\kappa_{ljk} \upsilon_{\nu_m^c}\upsilon_{\nu_j^c}\upsilon_{\nu_k^c} + 2Y_{\nu_{jk}} \kappa_{ikl} \upsilon_u \upsilon_{\nu_j} \upsilon_{\nu_l^c} \\
&\quad - Y_{\nu_{ji}} \lambda_k \upsilon_{\nu_j} \upsilon_{\nu_k^c} \upsilon_d - Y_{\nu_{kj}} \lambda_i \upsilon_{\nu_k} \upsilon_{\nu_j^c} \upsilon_d + Y_{\nu_{ji}} Y_{\nu_{lk}} \upsilon_{\nu_j} \upsilon_{\nu_l} \upsilon_{\nu_k^c} \\
&\quad + Y_{\nu_{ki}} Y_{\nu_{kj}} \upsilon_u^2 \upsilon_{\nu_j^c} = 0\,,
\end{aligned} \tag{A.4}$$

where $G^2 = g_1^2 + g_2^2$ and $g_1 c_{_W} = g_2 s_{_W} = e$.

Appendix B. Mass Matrices

In this appendix, we give the mass matrices in the $\mu\nu$SSM.

Appendix B.1. Scalar mass matrices

For this subsection, we use the indices $i, j, k, l, m = 1, 2, 3$ and $\alpha = 1, \ldots, 8$.

Appendix B.1.1. CP-even neutral scalars

In the unrotated basis $S'^T = (h_d, h_u, (\tilde{\nu}_i)^R, (\tilde{\nu}_i^c)^R)$, one can obtain the quadratic potential

$$V_{quadratic} = \frac{1}{2} S'^T M_S^2 S' \; . \tag{B.1}$$

And the expression for the independent coefficients of M_S^2 are given in detail below:

$$M^2_{h_d h_d} = m^2_{H_d} + \frac{G^2}{4}(3\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}) + \lambda_i \lambda_j \upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + \lambda_i\lambda_i\upsilon_u^2 \, , \tag{B.2}$$

$$\begin{aligned} M^2_{h_u h_u} = {} & m^2_{H_u} - \frac{G^2}{4}(\upsilon_d^2 - 3\upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}) + \lambda_i\lambda_j\upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + \lambda_i\lambda_i\upsilon_d^2 \\ & - 2Y_{\nu_{ij}}\lambda_j\upsilon_d\upsilon_{\nu_i} + Y_{\nu_{ki}}Y_{\nu_{kj}}\upsilon_{\nu_i^c}\upsilon_{\nu_j^c} + Y_{\nu_{ik}}Y_{\nu_{jk}}\upsilon_{\nu_i}\upsilon_{\nu_j} \, , \end{aligned} \tag{B.3}$$

$$\begin{aligned} M^2_{h_d h_u} = {} & -(A_\lambda \lambda)_i \upsilon_{\nu_i^c} - \frac{G^2}{2}\upsilon_d\upsilon_u + 2\lambda_i\lambda_i\upsilon_d\upsilon_u - \lambda_k\kappa_{ijk}\upsilon_{\nu_i^c}\upsilon_{\nu_j^c} \\ & - 2Y_{\nu_{ij}}\lambda_j\upsilon_u\upsilon_{\nu_i} \, , \end{aligned} \tag{B.4}$$

$$M^2_{h_d(\tilde{\nu}_i)^R} = \frac{G^2}{2}\upsilon_d\upsilon_{\nu_i} - Y_{\nu_{ij}}(\lambda_j\upsilon_u^2 + \lambda_k\upsilon_{\nu_k^c}\upsilon_{\nu_j^c}) \, , \tag{B.5}$$

$$\begin{aligned} M^2_{h_u(\tilde{\nu}_i)^R} = {} & -\frac{G^2}{2}\upsilon_u\upsilon_{\nu_i} + (A_\nu Y_\nu)_{ij}\upsilon_{\nu_j^c} - 2Y_{\nu_{ij}}\lambda_j\upsilon_d\upsilon_u + Y_{\nu_{ik}}\kappa_{ljk}\upsilon_{\nu_l^c}\upsilon_{\nu_j^c} \\ & + 2Y_{\nu_{ij}}Y_{\nu_{kj}}\upsilon_u\upsilon_{\nu_k} \, , \end{aligned} \tag{B.6}$$

$$\begin{aligned} M^2_{h_d(\tilde{\nu}_i^c)^R} = {} & -(A_\lambda\lambda)_i\upsilon_u + 2\lambda_i\lambda_j\upsilon_d\upsilon_{\nu_j^c} - 2\lambda_k\kappa_{ijk}\upsilon_u\upsilon_{\nu_j^c} \\ & - (Y_{\nu_{ji}}\lambda_k + Y_{\nu_{jk}}\lambda_i)\upsilon_{\nu_j}\upsilon_{\nu_k^c} \, , \end{aligned} \tag{B.7}$$

$$\begin{aligned} M^2_{h_u(\tilde{\nu}_i^c)^R} = {} & -(A_\lambda\lambda)_i\upsilon_d + (A_\nu Y_\nu)_{ji}\upsilon_{\nu_j} + 2\lambda_i\lambda_j\upsilon_u\upsilon_{\nu_j^c} - 2\lambda_k\kappa_{ijk}\upsilon_d\upsilon_{\nu_j^c} \\ & + 2Y_{\nu_{jk}}\kappa_{ilk}\upsilon_{\nu_j}\upsilon_{\nu_l^c} + 2Y_{\nu_{jk}}Y_{\nu_{ji}}\upsilon_u\upsilon_{\nu_k^c} \, , \end{aligned} \tag{B.8}$$

$$\begin{aligned} M^2_{(\tilde{\nu}_i)^R(\tilde{\nu}_j)^R} = {} & m^2_{\tilde{L}_{ij}} + \frac{G^2}{2}\upsilon_{\nu_i}\upsilon_{\nu_j} + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_k}\upsilon_{\nu_k})\delta_{ij} \\ & + Y_{\nu_{ik}}Y_{\nu_{jk}}\upsilon_u^2 + Y_{\nu_{ik}}Y_{\nu_{jl}}\upsilon_{\nu_k^c}\upsilon_{\nu_l^c} \, , \end{aligned} \tag{B.9}$$

$$\begin{aligned}
M^2_{(\tilde{\nu}_i)^R(\tilde{\nu}^c_j)^R} &= (A_\nu Y_\nu)_{ij}\upsilon_u - (Y_{\nu_{ij}}\lambda_k + Y_{\nu_{ik}}\lambda_j)\upsilon_d\upsilon_{\nu^c_k} + 2Y_{\nu_{ik}}\kappa_{jlk}\upsilon_u\upsilon_{\nu^c_l} \\
&\quad + (Y_{\nu_{ij}}Y_{\nu_{kl}} + Y_{\nu_{il}}Y_{\nu_{kj}})\upsilon_{\nu_k}\upsilon_{\nu^c_l}\ , \qquad \text{(B.10)} \\
M^2_{(\tilde{\nu}^c_i)^R(\tilde{\nu}^c_j)^R} &= m^2_{\tilde{\nu}^c_{ij}} + 2(A_\kappa\kappa)_{ijk}\upsilon_{\nu^c_k} - 2\lambda_k\kappa_{ijk}\upsilon_d\upsilon_u + \lambda_i\lambda_j(\upsilon_d^2 + \upsilon_u^2) \\
&\quad + (2\kappa_{ijk}\kappa_{lmk} + 4\kappa_{ilk}\kappa_{jmk})\upsilon_{\nu^c_l}\upsilon_{\nu^c_m} + 2Y_{\nu_{lk}}\kappa_{ijk}\upsilon_u\upsilon_{\nu_l} \\
&\quad - (Y_{\nu_{kj}}\lambda_i + Y_{\nu_{ki}}\lambda_j)\upsilon_d\upsilon_{\nu_k} + Y_{\nu_{ki}}(Y_{\nu_{kj}}\upsilon_u^2 + Y_{\nu_{lj}}\upsilon_{\nu_k}\upsilon_{\nu_l})\ . \qquad \text{(B.11)}
\end{aligned}$$

We can use an 8×8 unitary matrix R_S to diagonalize the mass matrix M_S^2

$$R_S^T M_S^2 R_S = (M_S^{diag})^2\ . \qquad \text{(B.12)}$$

By unitary matrix R_S, S'_α can be rotated to the mass eigenvectors S_α:

$$h_d = R_S^{1\alpha}S_\alpha,\ h_u = R_S^{2\alpha}S_\alpha,\ (\tilde{\nu}_i)^R = R_S^{(2+i)\alpha}S_\alpha,\ (\tilde{\nu}^c_i)^R = R_S^{(5+i)\alpha}S_\alpha\ . \qquad \text{(B.13)}$$

Appendix B.1.2. CP-odd neutral scalars

In the unrotated basis $P'^T = (P_d, P_u, (\tilde{\nu}_i)^I, (\tilde{\nu}^c_i)^I)$, one can also give the quadratic potential

$$V_{quadratic} = \frac{1}{2}P'^T M_P^2 P'\ , \qquad \text{(B.14)}$$

and the concrete expression for the independent coefficients of M_P^2

$$\begin{aligned}
M^2_{P_dP_d} &= m^2_{H_d} + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}) + \lambda_i\lambda_j\upsilon_{\nu^c_i}\upsilon_{\nu^c_j} + \lambda_i\lambda_i\upsilon_u^2\ , \qquad \text{(B.15)} \\
M^2_{P_uP_u} &= m^2_{H_u} - \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i}\upsilon_{\nu_i}) + \lambda_i\lambda_j\upsilon_{\nu^c_i}\upsilon_{\nu^c_j} + \lambda_i\lambda_i\upsilon_d^2 \\
&\quad - 2Y_{\nu_{ij}}\lambda_j\upsilon_d\upsilon_{\nu_i} + Y_{\nu_{ki}}Y_{\nu_{kj}}\upsilon_{\nu^c_i}\upsilon_{\nu^c_j} + Y_{\nu_{ik}}Y_{\nu_{jk}}\upsilon_{\nu_i}\upsilon_{\nu_j}\ , \qquad \text{(B.16)} \\
M^2_{P_dP_u} &= (A_\lambda\lambda)_i\upsilon_{\nu^c_i} + \lambda_k\kappa_{ijk}\upsilon_{\nu^c_i}\upsilon_{\nu^c_j}\ , \qquad \text{(B.17)} \\
M^2_{P_d(\tilde{\nu}_i)^I} &= -Y_{\nu_{ij}}(\lambda_j\upsilon_u^2 + \lambda_k\upsilon_{\nu^c_k}\upsilon_{\nu^c_j})\ , \qquad \text{(B.18)} \\
M^2_{P_u(\tilde{\nu}_i)^I} &= -(A_\nu Y_\nu)_{ij}\upsilon_{\nu^c_j} - Y_{\nu_{ik}}\kappa_{ljk}\upsilon_{\nu^c_l}\upsilon_{\nu^c_j}\ , \qquad \text{(B.19)} \\
M^2_{P_d(\tilde{\nu}^c_i)^I} &= (A_\lambda\lambda)_i\upsilon_u - 2\lambda_k\kappa_{ijk}\upsilon_u\upsilon_{\nu^c_j} - (Y_{\nu_{ji}}\lambda_k - Y_{\nu_{jk}}\lambda_i)\upsilon_{\nu_j}\upsilon_{\nu^c_k}\ , \qquad \text{(B.20)} \\
M^2_{P_u(\tilde{\nu}^c_i)^I} &= (A_\lambda\lambda)_i\upsilon_d - (A_\nu Y_\nu)_{ji}\upsilon_{\nu_j} - 2(\lambda_k\kappa_{ilk}\upsilon_d - Y_{\nu_{jk}}\kappa_{ilk}\upsilon_{\nu_j})\upsilon_{\nu^c_l}\ , \qquad \text{(B.21)} \\
M^2_{(\tilde{\nu}_i)^I(\tilde{\nu}_j)^I} &= m^2_{\tilde{L}_{ij}} + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_k}\upsilon_{\nu_k})\delta_{ij} + Y_{\nu_{ik}}Y_{\nu_{jk}}\upsilon_u^2
\end{aligned}$$

$$\begin{aligned}
&\quad + Y_{\nu_{ik}} Y_{\nu_{jl}} \upsilon_{\nu^c_k} \upsilon_{\nu^c_l} \;, && \text{(B.22)}\\
M^2_{(\tilde{\nu}_i)^I (\tilde{\nu}^c_j)^I} &= -(A_\nu Y_\nu)_{ij} \upsilon_u + (Y_{\nu_{ij}} \lambda_k - Y_{\nu_{ik}} \lambda_j) \upsilon_d \upsilon_{\nu^c_k} + 2 Y_{\nu_{il}} \kappa_{jlk} \upsilon_u \upsilon_{\nu^c_k} \\
&\quad - (Y_{\nu_{ij}} Y_{\nu_{kl}} - Y_{\nu_{il}} Y_{\nu_{kj}}) \upsilon_{\nu_k} \upsilon_{\nu^c_l} \;, && \text{(B.23)}\\
M^2_{(\tilde{\nu}^c_i)^I (\tilde{\nu}^c_j)^I} &= m^2_{\tilde{\nu}^c_{ij}} - 2(A_\kappa \kappa)_{ijk} \upsilon_{\nu^c_k} + 2\lambda_k \kappa_{ijk} \upsilon_d \upsilon_u + \lambda_i \lambda_j (\upsilon_d^2 + \upsilon_u^2) \\
&\quad - (2\kappa_{ijk} \kappa_{lmk} - 4\kappa_{imk} \kappa_{ljk}) \upsilon_{\nu^c_l} \upsilon_{\nu^c_m} - 2 Y_{\nu_{lk}} \kappa_{ijk} \upsilon_u \upsilon_{\nu_l} \\
&\quad - (Y_{\nu_{kj}} \lambda_i + Y_{\nu_{ki}} \lambda_j) \upsilon_d \upsilon_{\nu_k} + Y_{\nu_{ki}} (Y_{\nu_{kj}} \upsilon_u^2 + Y_{\nu_{lj}} \upsilon_{\nu_k} \upsilon_{\nu_l}) \;. && \text{(B.24)}
\end{aligned}$$

Using an 8×8 unitary matrix R_P to diagonalize the mass matrix M_P^2

$$R_P^T M_P^2 R_P = (M_P^{diag})^2 \;, \qquad \text{(B.25)}$$

we can obtain the mass eigenvectors P_α:

$$P_d = R_P^{1\alpha} P_\alpha,\; P_u = R_P^{2\alpha} P_\alpha,\; (\tilde{\nu}_i)^I = R_P^{(2+i)\alpha} P_\alpha,\; (\tilde{\nu}^c_i)^I = R_P^{(5+i)\alpha} P_\alpha \;. \qquad \text{(B.26)}$$

Appendix B.1.3. Charged scalars

The quadratic potential includes

$$V_{quadratic} = S'^{-T} M^2_{S^\pm} S'^{+} \;, \qquad \text{(B.27)}$$

where $S'^{\pm T} = (H_d^\pm, H_u^\pm, \tilde{e}^\pm_{L_i}, \tilde{e}^\pm_{R_i})$ is in the unrotated basis, $\tilde{e}^-_{L_i} \equiv \tilde{e}_i$ and $\tilde{e}^+_{R_i} \equiv \tilde{e}^c_i$. The concrete expression for the independent coefficients of $M^2_{S^\pm}$ are given below:

$$\begin{aligned}
M^2_{H_d^\pm H_d^\pm} &= m^2_{H_d} + \frac{g_2^2}{2}(\upsilon_u^2 - \upsilon_{\nu_i} \upsilon_{\nu_i}) + \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i} \upsilon_{\nu_i}) + \lambda_i \lambda_j \upsilon_{\nu^c_i} \upsilon_{\nu^c_j} \\
&\quad + Y_{e_{ik}} Y_{e_{jk}} \upsilon_{\nu_i} \upsilon_{\nu_j} \;, && \text{(B.28)}\\
M^2_{H_u^\pm H_u^\pm} &= m^2_{H_u} + \frac{g_2^2}{2}(\upsilon_d^2 + \upsilon_{\nu_i} \upsilon_{\nu_i}) - \frac{G^2}{4}(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_i} \upsilon_{\nu_i}) + \lambda_i \lambda_j \upsilon_{\nu^c_i} \upsilon_{\nu^c_j} \\
&\quad + Y_{\nu_{ik}} Y_{\nu_{ij}} \upsilon_{\nu^c_j} \upsilon_{\nu^c_k} \;, && \text{(B.29)}\\
M^2_{H_d^\pm H_u^\pm} &= (A_\lambda \lambda)_i \upsilon_{\nu^c_i} + \frac{g_2^2}{2} \upsilon_d \upsilon_u - \lambda_i \lambda_i \upsilon_d \upsilon_u + \lambda_k \kappa_{ijk} \upsilon_{\nu^c_i} \upsilon_{\nu^c_j} \\
&\quad + Y_{\nu_{ij}} \lambda_j \upsilon_u \upsilon_{\nu_i} \;, && \text{(B.30)}\\
M^2_{H_d^\pm \tilde{e}^\pm_{L_i}} &= \frac{g_2^2}{2} \upsilon_d \upsilon_{\nu_i} - Y_{\nu_{ij}} \lambda_k \upsilon_{\nu^c_k} \upsilon_{\nu^c_j} - Y_{e_{ij}} Y_{e_{kj}} \upsilon_d \upsilon_{\nu_k} \;, && \text{(B.31)}\\
M^2_{H_u^\pm \tilde{e}^\pm_{L_i}} &= \frac{g_2^2}{2} \upsilon_u \upsilon_{\nu_i} - (A_\nu Y_\nu)_{ij} \upsilon_{\nu^c_j} + Y_{\nu_{ij}} \lambda_j \upsilon_d \upsilon_u - Y_{\nu_{ij}} \kappa_{ljk} \upsilon_{\nu^c_l} \upsilon_{\nu^c_k}
\end{aligned}$$

$$- Y_{\nu_{ik}} Y_{\nu_{kj}} \upsilon_u \upsilon_{\nu_j} \ , \tag{B.32}$$

$$M^2_{H_d^\pm \tilde{e}^\pm_{R_i}} = -(A_e Y_e)_{ji} \upsilon_{\nu_j} - Y_{e_{ki}} Y_{\nu_{kj}} \upsilon_u \upsilon_{\nu^c_j} \ , \tag{B.33}$$

$$M^2_{H_u^\pm \tilde{e}^\pm_{R_i}} = -Y_{e_{ki}} (\lambda_j \upsilon_{\nu^c_j} \upsilon_{\nu_k} + Y_{\nu_{kj}} \upsilon_d \upsilon_{\nu^c_j}) \ , \tag{B.34}$$

$$\begin{aligned} M^2_{\tilde{e}^\pm_{L_i} \tilde{e}^\pm_{L_j}} &= m^2_{\tilde{L}_{ij}} + \frac{1}{4}(g_1^2 - g_2^2)(\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_k} \upsilon_{\nu_k}) \delta_{ij} + \frac{g_2^2}{2} \upsilon_{\nu_i} \upsilon_{\nu_j} \\ &\quad + Y_{\nu_{il}} Y_{\nu_{jk}} \upsilon_{\nu^c_l} \upsilon_{\nu^c_k} + Y_{e_{ik}} Y_{e_{jk}} \upsilon_d^2 \ , \end{aligned} \tag{B.35}$$

$$M^2_{\tilde{e}^\pm_{L_i} \tilde{e}^\pm_{R_j}} = (A_e Y_e)_{ij} \upsilon_d - Y_{e_{ij}} \lambda_k \upsilon_u \upsilon_{\nu^c_k} \ , \tag{B.36}$$

$$\begin{aligned} M^2_{\tilde{e}^\pm_{R_i} \tilde{e}^\pm_{R_j}} &= m^2_{\tilde{e}^c_{ij}} - \frac{1}{2} g_1^2 (\upsilon_d^2 - \upsilon_u^2 + \upsilon_{\nu_k} \upsilon_{\nu_k}) \delta_{ij} + Y_{e_{ki}} Y_{e_{kj}} \upsilon_d^2 \\ &\quad + Y_{e_{li}} Y_{e_{kj}} \upsilon_{\nu_k} \upsilon_{\nu_l} \ . \end{aligned} \tag{B.37}$$

Through an 8 × 8 unitary matrix $R_{S^\pm}$ to diagonalize the mass matrix $M^2_{S^\pm}$

$$R^T_{S^\pm} M^2_{S^\pm} R_{S^\pm} = (M^{diag}_{S^\pm})^2 \ , \tag{B.38}$$

$S'^\pm_\alpha$ can be rotated to the mass eigenvectors $S^\pm_\alpha$:

$$H_d^\pm = R^{1\alpha}_{S^\pm} S^\pm_\alpha, \ H_u^\pm = R^{2\alpha}_{S^\pm} S^\pm_\alpha, \ \tilde{e}^\pm_{L_i} = R^{(2+i)\alpha}_{S^\pm} S^\pm_\alpha, \ \tilde{e}^\pm_{R_i} = R^{(5+i)\alpha}_{S^\pm} S^\pm_\alpha \ . \tag{B.39}$$

Appendix B.2. Neutral fermion mass matrix

Neutrinos mix with the neutralinos and therefore in the unrotated basis $\chi'^{\circ T} = \Big(\tilde{B}^\circ, \tilde{W}^\circ, \tilde{H}_d, \tilde{H}_u, \nu_{R_i}, \nu_{L_i} \Big)$, one can have the neutral fermion mass terms in the Lagrangian:

$$- \frac{1}{2} \chi'^{\circ T} M_n \chi'^\circ + \text{H.c.} \ , \tag{B.40}$$

where

$$M_n = \begin{pmatrix} M & m^T \\ m & 0_{3\times 3} \end{pmatrix} , \tag{B.41}$$

with

$$m = \begin{pmatrix} -\frac{g_1}{\sqrt{2}} \upsilon_{\nu_1} & \frac{g_2}{\sqrt{2}} \upsilon_{\nu_1} & 0 & Y_{\nu_{1i}} \upsilon_{\nu^c_i} & Y_{\nu_{11}} \upsilon_u & Y_{\nu_{12}} \upsilon_u & Y_{\nu_{13}} \upsilon_u \\ -\frac{g_1}{\sqrt{2}} \upsilon_{\nu_2} & \frac{g_2}{\sqrt{2}} \upsilon_{\nu_2} & 0 & Y_{\nu_{2i}} \upsilon_{\nu^c_i} & Y_{\nu_{21}} \upsilon_u & Y_{\nu_{22}} \upsilon_u & Y_{\nu_{23}} \upsilon_u \\ -\frac{g_1}{\sqrt{2}} \upsilon_{\nu_3} & \frac{g_2}{\sqrt{2}} \upsilon_{\nu_3} & 0 & Y_{\nu_{3i}} \upsilon_{\nu^c_i} & Y_{\nu_{31}} \upsilon_u & Y_{\nu_{32}} \upsilon_u & Y_{\nu_{33}} \upsilon_u \end{pmatrix} \tag{B.42}$$

and

$$M = \begin{pmatrix} M_1 & 0 & \frac{-g_1}{\sqrt{2}}\upsilon_d & \frac{g_1}{\sqrt{2}}\upsilon_u & 0 & 0 & 0 \\ 0 & M_2 & \frac{g_2}{\sqrt{2}}\upsilon_d & \frac{-g_2}{\sqrt{2}}\upsilon_u & 0 & 0 & 0 \\ \frac{-g_1}{\sqrt{2}}\upsilon_d & \frac{g_2}{\sqrt{2}}\upsilon_d & 0 & -\lambda_i\upsilon_{\nu^c_i} & -\lambda_1\upsilon_u & -\lambda_2\upsilon_u & -\lambda_3\upsilon_u \\ \frac{g_1}{\sqrt{2}}\upsilon_u & \frac{-g_2}{\sqrt{2}}\upsilon_u & -\lambda_i\upsilon_{\nu^c_i} & 0 & y_1 & y_2 & y_3 \\ 0 & 0 & -\lambda_1\upsilon_u & y_1 & 2\kappa_{11j}\upsilon_{\nu^c_j} & 2\kappa_{12j}\upsilon_{\nu^c_j} & 2\kappa_{13j}\upsilon_{\nu^c_j} \\ 0 & 0 & -\lambda_2\upsilon_u & y_2 & 2\kappa_{21j}\upsilon_{\nu^c_j} & 2\kappa_{22j}\upsilon_{\nu^c_j} & 2\kappa_{23j}\upsilon_{\nu^c_j} \\ 0 & 0 & -\lambda_3\upsilon_u & y_3 & 2\kappa_{31j}\upsilon_{\nu^c_j} & 2\kappa_{32j}\upsilon_{\nu^c_j} & 2\kappa_{33j}\upsilon_{\nu^c_j} \end{pmatrix} \tag{B.43}$$

where $y_i = -\lambda_i\upsilon_d + Y_{\nu_{ji}}\upsilon_{\nu_j}$. Here, the submatrix m is neutralino-neutrino mixing, and the submatrix M is neutralino mass matrix. This 10×10 symmetric matrix M_n can be diagonalized by a 10×10 unitary matrix Z_n:

$$Z_n^T M_n Z_n = M_{nd} \,, \tag{B.44}$$

where M_{nd} is the diagonal neutral fermion mass matrix. Then, we have the neutral fermion mass eigenstates:

$$\chi^\circ_\alpha = \left(\frac{\kappa^\circ_\alpha}{\kappa^\circ_\alpha} \right), \quad \alpha = 1, \ldots, 10 \tag{B.45}$$

with

$$\begin{cases} \tilde{B}^\circ = Z_n^{1\alpha}\kappa^\circ_\alpha \,, & \tilde{H}_d = Z_n^{3\alpha}\kappa^\circ_\alpha \,, \quad \nu_{R_i} = Z_n^{(4+i)\alpha}\kappa^\circ_\alpha \,, \\ \tilde{W}^\circ = Z_n^{2\alpha}\kappa^\circ_\alpha \,, & \tilde{H}_u = Z_n^{4\alpha}\kappa^\circ_\alpha \,, \quad \nu_{L_i} = Z_n^{(7+i)\alpha}\kappa^\circ_\alpha \,. \end{cases} \tag{B.46}$$

Appendix B.3. Charged fermion mass matrix

Charged leptons mix with the charginos and therefore in the unrotated basis where $\Psi^{-T} = \left(-i\tilde{\lambda}^-, \tilde{H}_d^-, e_{L_i}^-\right)$ and $\Psi^{+T} = \left(-i\tilde{\lambda}^+, \tilde{H}_u^+, e_{R_i}^+\right)$, one can obtain the charged fermion mass terms in the Lagrangian:

$$-\Psi^{-T} M_c \Psi^+ + \text{H.c.} \,, \tag{B.47}$$

where

$$M_c = \begin{pmatrix} M_\pm & b \\ c & m_l \end{pmatrix} . \tag{B.48}$$

Here, the submatrix $M_\pm$ is chargino mass matrix

$$M_\pm = \left(\begin{array}{cc} M_2 & g_2 \upsilon_u \\ g_2 \upsilon_d & \lambda_i \upsilon_{\nu_i^c} \end{array} \right) . \tag{B.49}$$

And the submatrices b and c give rise to chargino-charged lepton mixing. They are defined as

$$b = \left(\begin{array}{ccc} 0 & 0 & 0 \\ -Y_{e_{i1}} \upsilon_{\nu_i} & -Y_{e_{i2}} \upsilon_{\nu_i} & -Y_{e_{i3}} \upsilon_{\nu_i} \end{array} \right) , \tag{B.50}$$

$$c = \left(\begin{array}{cc} g_2 \upsilon_{\nu_1} & -Y_{\nu_{1i}} \upsilon_{\nu_i^c} \\ g_2 \upsilon_{\nu_2} & -Y_{\nu_{2i}} \upsilon_{\nu_i^c} \\ g_2 \upsilon_{\nu_3} & -Y_{\nu_{3i}} \upsilon_{\nu_i^c} \end{array} \right) . \tag{B.51}$$

And the submatrix m_l is the charged lepton mass matrix

$$m_l = \left(\begin{array}{ccc} Y_{e_{11}} \upsilon_d & Y_{e_{12}} \upsilon_d & Y_{e_{13}} \upsilon_d \\ Y_{e_{21}} \upsilon_d & Y_{e_{22}} \upsilon_d & Y_{e_{23}} \upsilon_d \\ Y_{e_{31}} \upsilon_d & Y_{e_{32}} \upsilon_d & Y_{e_{33}} \upsilon_d \end{array} \right) . \tag{B.52}$$

This 5×5 mass matrix M_c can be diagonalized by the 5×5 unitary matrices Z_- and Z_+:

$$Z_-^T M_c Z_+ = M_{cd} \, , \tag{B.53}$$

where M_{cd} is the diagonal charged fermion mass matrix. Then, one can obtain the charged fermion mass eigenstates:

$$\chi_\alpha = \left(\begin{array}{c} \kappa_\alpha^- \\ \kappa_\alpha^+ \end{array} \right) , \quad \alpha = 1, \ldots, 5 \tag{B.54}$$

with

$$\left\{ \begin{array}{l} \tilde{\lambda}^- = i Z_-^{1\alpha} \kappa_\alpha^- \, , \quad \tilde{H}_d^- = Z_-^{2\alpha} \kappa_\alpha^- \, , \quad e_{L_i} = Z_-^{(2+i)\alpha} \kappa_\alpha^- ; \\ \tilde{\lambda}^+ = i Z_+^{1\alpha} \kappa_\alpha^+ \, , \quad \tilde{H}_u^+ = Z_+^{2\alpha} \kappa_\alpha^+ \, , \quad e_{R_i} = Z_+^{(2+i)\alpha} \kappa_\alpha^+ \, . \end{array} \right. \tag{B.55}$$

Appendix C. Approximate diagonalization of mass matrices

Appendix C.1. Neutral fermion mass matrix

If the R-parity breaking parameters are small in the sense that for [16, 28]

$$\xi = m.M^{-1}, \tag{C.1}$$

all $\xi_{ij} \ll 1$, one can find an approximate diagonalization of neutral fermion mass matrix. In leading order in ξ, the rotation matrix Z_n is given by

$$Z_n = \begin{pmatrix} 1-\frac{1}{2}\xi^T\xi & -\xi^T \\ \xi & 1-\frac{1}{2}\xi\xi^T \end{pmatrix} \begin{pmatrix} V & 0 \\ 0 & U_\nu \end{pmatrix}. \tag{C.2}$$

The first matrix in (C.2) above approximately block-diagonalizes the matrix M_n to the form $diag\,(M, m_{eff})$, where

$$m_{eff} = -m.M^{-1}.m^T\ . \tag{C.3}$$

The submatrices V and U_ν respectively diagonalize M and m_{eff} in the following way:

$$\left\{ \begin{array}{l} V^T M V = M_d\ , \\ U_\nu^T m_{eff} U_\nu = m_{\nu d}\ , \end{array} \right. \tag{C.4}$$

where M_d and $m_{\nu d}$ are respectively diagonal neutralino and neutrino mass matrix.

Appendix C.2. Charged fermion mass matrix

Similarly to the approximate diagonalization of the neutral fermion mass matrix discussed above, it's also possible to find an approximate diagonalization procedure of the charged fermion mass matrix for the small R-parity breaking parameters [28]. Then, we can define

$$\left\{ \begin{array}{l} \xi_L = c.M_\pm^{-1} + m_l.b^T.(M_\pm^{-1})^T.M_\pm^{-1}; \\ \xi_R = b^T.(M_\pm^{-1})^T + {m_l}^T.c.M_\pm^{-1}.(M_\pm^{-1})^T. \end{array} \right. \tag{C.5}$$

All $\xi_{L_{ij}} \ll 1$ and $\xi_{R_{ij}} \ll 1$, so in leading order in ξ_L and ξ_R, the rotation matrices Z_- and Z_+ are respectively given by

$$Z_- = \begin{pmatrix} 1-\frac{1}{2}\xi_L^T\xi_L & -\xi_L^T \\ \xi_L & 1-\frac{1}{2}\xi_L\xi_L^T \end{pmatrix} \begin{pmatrix} U_- & 0 \\ 0 & V_- \end{pmatrix}, \tag{C.6}$$

$$Z_+ = \left(\begin{array}{cc} 1-\frac{1}{2}\xi_R^T\xi_R & -\xi_R^T \\ \xi_R & 1-\frac{1}{2}\xi_R\xi_R^T \end{array} \right) \left(\begin{array}{cc} U_+ & 0 \\ 0 & V_+ \end{array} \right). \tag{C.7}$$

Then the matrix M_c can approximately be block-diagonalized to the form $diag\,(M_\pm, m_l)$. And the submatrices U_-, U_+ and V_-, V_+ respectively diagonalize $M_\pm$ and m_l in the following way:

$$\left\{ \begin{array}{l} U_-^T M_\pm U_+ = M_{\pm d}\ , \\ V_-^T m_l V_+ = m_{ld}\ , \end{array} \right. \tag{C.8}$$

where $M_{\pm d}$ and m_{ld} are respectively diagonal chargino and charged lepton mass matrix.

Appendix D. Interaction Lagrangian

In this part, we give the interaction Lagrangian of the relative vertices for the LFV processes in the $\mu\nu$SSM. And we use the indices $i,j=1,\ldots,3$, $\beta,\zeta=1,\ldots,5$, $\alpha,\rho=1,\ldots,8$ and $\eta=1,\ldots,10$.

Appendix D.1. Charged fermion-neutral fermion-gauge boson

We now give the interaction Lagrangian of charged fermion, neutral fermion and gauge boson,

$$\begin{aligned} \mathcal{L}_{int} = & eF_\mu\bar{\chi}_\beta\gamma^\mu\chi_\beta + Z_\mu\bar{\chi}_\beta(C_L^{Z\chi_\zeta\bar{\chi}_\beta}\gamma^\mu P_L + C_R^{Z\chi_\zeta\bar{\chi}_\beta}\gamma^\mu P_R)\chi_\zeta \\ & + W_\mu^+\bar{\chi}_\eta^0(C_L^{W\chi_\beta\bar{\chi}_\eta^0}\gamma^\mu P_L + C_R^{W\chi_\beta\bar{\chi}_\eta^0}\gamma^\mu P_R)\chi_\beta \\ & + W_\mu^-\bar{\chi}_\beta(C_L^{W\chi_\eta^0\bar{\chi}_\beta}\gamma^\mu P_L + C_R^{W\chi_\eta^0\bar{\chi}_\beta}\gamma^\mu P_R)\chi_\eta^0 + \cdots, \end{aligned} \tag{D.1}$$

where the coefficients are

$$\begin{aligned} C_L^{Z\chi_\zeta\bar{\chi}_\beta} &= \frac{e}{2s_{_W}c_{_W}}\Big[(1-2s_{_W}^2)\delta^{\zeta\beta} + Z_-^{1\zeta*}Z_-^{1\beta}\Big]\ , \\ C_R^{Z\chi_\zeta\bar{\chi}_\beta} &= \frac{e}{2s_{_W}c_{_W}}\Big[2Z_+^{1\zeta*}Z_+^{1\beta} + Z_+^{2\zeta*}Z_+^{2\beta} - 2s_{_W}^2\delta^{\zeta\beta}\Big]\ , \\ C_L^{W\chi_\beta\bar{\chi}_\eta^\circ} &= -\frac{e}{\sqrt{2}s_{_W}}\Big[\sqrt{2}Z_-^{1\beta}Z_n^{2\eta*} + Z_-^{2\beta}Z_n^{3\eta*} + Z_-^{(2+i)\beta}Z_n^{(7+i)\eta*}\Big]\ , \\ C_R^{W\chi_\beta\bar{\chi}_\eta^\circ} &= -\frac{e}{\sqrt{2}s_{_W}}\Big[\sqrt{2}Z_+^{1\beta*}Z_n^{2\eta} - Z_+^{2\beta*}Z_n^{4\eta}\Big]\ , \\ C_L^{W\chi_\eta^\circ\bar{\chi}_\beta} &= \Big[C_L^{W\chi_\beta\bar{\chi}_\eta^\circ}\Big]^*, \qquad C_R^{W\chi_\eta^\circ\bar{\chi}_\beta} = \Big[C_R^{W\chi_\beta\bar{\chi}_\eta^\circ}\Big]^*. \end{aligned} \tag{D.2}$$

Appendix D.2. Charged scalars-gauge boson

The interaction Lagrangian of charged scalars and gauge boson is written as

$$\mathcal{L}_{int} = ieF_\mu S_\alpha^{-*} \overleftrightarrow{\partial^\mu} S_\alpha^- + ieC^{ZS_\alpha^- S_\rho^{-*}} Z_\mu S_\rho^{-*} \overleftrightarrow{\partial^\mu} S_\alpha^- + \cdots . \tag{D.3}$$

The coefficient is

$$C^{ZS_\alpha^- S_\rho^{-*}} = \frac{e}{2s_{_W} c_{_W}} \Big[(1-2s_{_W}^2)\delta^{\alpha\rho} - R_{S^\pm}^{(5+i)\alpha^*} R_{S^\pm}^{(5+i)\rho} \Big] . \tag{D.4}$$

Appendix D.3. Charged fermion-neutral fermion-scalars

The interaction Lagrangian of charged fermion, neutral fermion and scalars is similarly written by

$$\begin{aligned}\mathcal{L}_{int} = {} & S_\alpha \bar{\chi}_\zeta (C_L^{S_\alpha \chi_\beta \bar{\chi}_\zeta} P_L + C_R^{S_\alpha \chi_\beta \bar{\chi}_\zeta} P_R)\chi_\beta + P_\alpha \bar{\chi}_\zeta (C_L^{P_\alpha \chi_\beta \bar{\chi}_\zeta} P_L \\ & + C_R^{P_\alpha \chi_\beta \bar{\chi}_\zeta} P_R)\chi_\beta + S_\alpha^- \bar{\chi}_\beta (C_L^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_\beta} P_L + C_R^{S_\alpha^- \chi_\eta^\circ \bar{\chi}_\beta} P_R)\chi_\eta^0 \\ & + S_\alpha^{-*} \bar{\chi}_\eta^0 (C_L^{S_\alpha^{-*} \chi_\beta \bar{\chi}_\eta^\circ} P_L + C_R^{S_\alpha^{-*} \chi_\beta \bar{\chi}_\eta^\circ} P_R)\chi_\beta + \cdots . \end{aligned} \tag{D.5}$$

And the coefficients are

$$\begin{aligned} C_L^{S_\alpha \chi_\beta \bar{\chi}_\zeta} = {} & \frac{-e}{\sqrt{2}s_{_W}} \Big[R_S^{2\alpha} Z_-^{1\beta} Z_+^{2\zeta} + R_S^{1\alpha} Z_-^{2\beta} Z_+^{1\zeta} + R_S^{(5+i)\alpha} Z_-^{(2+i)\beta} Z_+^{1\zeta} \Big] \\ & + \frac{1}{\sqrt{2}} Y_{e_{ij}} \Big[R_S^{(5+i)\alpha} Z_-^{1\beta} Z_+^{(2+j)\zeta} - R_S^{1\alpha} Z_-^{(2+i)\beta} Z_+^{(2+j)\zeta} \Big] \\ & - \frac{1}{\sqrt{2}} Y_{\nu_{ij}} R_S^{(2+j)\alpha} Z_-^{(2+i)\beta} Z_+^{2\zeta} - \frac{1}{\sqrt{2}} \lambda_i R_S^{(2+i)\alpha} Z_-^{2\beta} Z_+^{2\zeta} , \\ C_L^{P_\alpha \chi_\beta \bar{\chi}_\zeta} = {} & \frac{ie}{\sqrt{2}s_{_W}} \Big[R_P^{2\alpha} Z_-^{1\beta} Z_+^{2\zeta} + R_P^{1\alpha} Z_-^{2\beta} Z_+^{1\zeta} + R_P^{(5+i)\alpha} Z_-^{(2+i)\beta} Z_+^{1\zeta} \Big] \\ & + \frac{i}{\sqrt{2}} Y_{e_{ij}} \Big[R_P^{(5+i)\alpha} Z_-^{1\beta} Z_+^{(2+j)\zeta} - R_P^{1\alpha} Z_-^{(2+i)\beta} Z_+^{(2+j)\zeta} \Big] \\ & - \frac{i}{\sqrt{2}} Y_{\nu_{ij}} R_P^{(2+j)\alpha} Z_-^{(2+i)\beta} Z_+^{2\zeta} - \frac{i}{\sqrt{2}} \lambda_i R_P^{(2+i)\alpha} Z_-^{2\beta} Z_+^{2\zeta} , \\ C_L^{S_\alpha^- \chi_\eta^0 \bar{\chi}_\beta} = {} & \frac{-e}{\sqrt{2}s_W c_W} R_{S^\pm}^{2\alpha *} Z_+^{2\beta} \Big[c_W Z_n^{2\eta} + s_W Z_n^{1\eta} \Big] - \frac{e}{s_W} R_{S^\pm}^{2\alpha *} Z_+^{1\beta} Z_n^{4\eta} \\ & - \frac{\sqrt{2}e}{s_W} R_{S^\pm}^{(5+i)\alpha *} Z_+^{(2+i)\beta} Z_n^{1\eta} + Y_{\nu_{ij}} R_{S^\pm}^{(2+i)\alpha} Z_+^{2\beta} Z_n^{(4+j)\eta} \end{aligned}$$

$$\begin{aligned}
&+Y_{e_{ij}}Z_{+}^{(2+j)\beta}\Big[R_{S^\pm}^{1\alpha}Z_n^{(7+i)\eta}-R_{S^\pm}^{(2+i)\alpha}Z_n^{3\eta}\Big]-\lambda_i R_{S^\pm}^{1\alpha}Z_{+}^{2\beta}Z_n^{(4+i)\eta},\\
C_L^{S_\alpha^{-*}\chi_\beta\bar{\chi}_\eta^\circ} &= \frac{e}{\sqrt{2}s_W c_W}\Big[R_{S^\pm}^{1\alpha*}Z_{-}^{2\beta}+R_{S^\pm}^{(2+i)\alpha^*}Z_{-}^{(2+i)\beta}\Big]\Big[c_W Z_n^{2\eta}+s_W Z_n^{1\eta}\Big]\\
&-\frac{e}{s_W}Z_{-}^{1\beta}\Big[R_{S^\pm}^{1\alpha*}Z_n^{3\eta}+R_{S^\pm}^{(2+i)\alpha*}Z_n^{(7+i)\eta}\Big]+Y_{\nu_{ij}}R_{S^\pm}^{2\alpha}Z_{-}^{(2+i)\beta}Z_n^{(4+j)\eta}\\
&+Y_{e_{ij}}R_{S^\pm}^{(5+j)\alpha}\Big[Z_{-}^{2\beta}Z_n^{(7+i)\eta}-Z_{-}^{(2+i)\beta}Z_n^{3\eta}\Big]-\lambda_i R_{S^\pm}^{2\alpha}Z_{-}^{2\beta}Z_n^{(4+i)\eta},\\
C_R^{S_\alpha\chi_\beta\bar{\chi}_\zeta} &= \Big[C_L^{S_\alpha\chi_\zeta\bar{\chi}_\beta}\Big]^*, \qquad C_R^{P_\alpha\chi_\beta\bar{\chi}_\zeta} = \Big[C_L^{P_\alpha\chi_\zeta\bar{\chi}_\beta}\Big]^*,\\
C_R^{S_\alpha^{-}\chi_\eta^\circ\bar{\chi}_\beta} &= \Big[C_L^{S_\alpha^{-*}\chi_\beta\bar{\chi}_\eta^0}\Big]^*, \qquad C_R^{S_\alpha^{-*}\chi_\beta\bar{\chi}_\eta^\circ} = \Big[C_L^{S_\alpha^{-}\chi_\eta^\circ\bar{\chi}_\beta}\Big]^*. \qquad \text{(D.6)}
\end{aligned}$$

Appendix E. Loop-momentum integral

Defining $x_i = \frac{m_i^2}{m_W^2}$, we can find the loop-momentum integral for $l_j^- \to l_i^- \gamma$:

$$I_1(x_1,x_2) = \frac{1}{16\pi^2}\Big[\frac{1+\ln x_2}{(x_2-x_1)}+\frac{x_1\ln x_1 - x_2\ln x_2}{(x_2-x_1)^2}\Big], \qquad \text{(E.1)}$$

$$I_2(x_1,x_2) = \frac{1}{16\pi^2}\Big[-\frac{1+\ln x_1}{(x_2-x_1)}-\frac{x_1\ln x_1 - x_2\ln x_2}{(x_2-x_1)^2}\Big], \qquad \text{(E.2)}$$

$$\begin{aligned}
I_3(x_1,x_2) &= \frac{1}{32\pi^2}\Big[\frac{3+2\ln x_2}{(x_2-x_1)}-\frac{2x_2+4x_2\ln x_2}{(x_2-x_1)^2}-\frac{2x_1^2\ln x_1}{(x_2-x_1)^3}\\
&+\frac{2x_2^2\ln x_2}{(x_2-x_1)^3}\Big], \qquad \text{(E.3)}
\end{aligned}$$

$$\begin{aligned}
I_4(x_1,x_2) &= \frac{1}{96\pi^2}\Big[\frac{11+6\ln x_2}{(x_2-x_1)}-\frac{15x_2+18x_2\ln x_2}{(x_2-x_1)^2}+\frac{6x_2^2+18x_2^2\ln x_2}{(x_2-x_1)^3}\\
&+\frac{6x_1^3\ln x_1-6x_2^3\ln x_2}{(x_2-x_1)^4}\Big]. \qquad \text{(E.4)}
\end{aligned}$$

And we also can find the loop-momentum integral for $l_j^- \to l_i^- l_i^- l_i^+$:

$$\begin{aligned}
G_1(x_1,x_2,x_3) &= \frac{1}{16\pi^2}\Big[\frac{x_1\ln x_1}{(x_1-x_2)(x_1-x_3)}+\frac{x_2\ln x_2}{(x_2-x_1)(x_2-x_3)}\\
&+\frac{x_3\ln x_3}{(x_3-x_1)(x_3-x_2)}\Big], \qquad \text{(E.5)}\\
G_2(x_1,x_2,x_3) &= \frac{1}{16\pi^2}\Big[-(\Delta+1+\ln x_\mu)+\frac{x_1^2\ln x_1}{(x_1-x_2)(x_1-x_3)}
\end{aligned}$$

$$+\frac{x_2^2\ln x_2}{(x_2-x_1)(x_2-x_3)}+\frac{x_3^2\ln x_3}{(x_3-x_1)(x_3-x_2)}\Big].\tag{E.6}$$

Here, $x_\mu = \frac{\mu^2}{m_W^2}$. $G_2(x_1,x_2,x_3)$ is divergence, so here we use dimensional regularization to cancel the divergent part $(\Delta+1+\ln x_\mu)$. In the numerical calculation, we will keep the remaining convergent part.

$$\begin{aligned}G_3(x_1,x_2,x_3,x_4) &= \frac{1}{16\pi^2}\Big[\frac{x_1\ln x_1}{(x_1-x_2)(x_1-x_3)(x_1-x_4)}\\ &+\frac{x_2\ln x_2}{(x_2-x_1)(x_2-x_3)(x_2-x_4)}+\frac{x_3\ln x_3}{(x_3-x_1)(x_3-x_2)(x_3-x_4)}\\ &+\frac{x_4\ln x_4}{(x_4-x_1)(x_4-x_2)(x_4-x_3)}\Big],\end{aligned}\tag{E.7}$$

$$\begin{aligned}G_4(x_1,x_2,x_3,x_4) &= \frac{1}{16\pi^2}\Big[\frac{x_1^2\ln x_1}{(x_1-x_2)(x_1-x_3)(x_1-x_4)}\\ &+\frac{x_2^2\ln x_2}{(x_2-x_1)(x_2-x_3)(x_2-x_4)}+\frac{x_3^2\ln x_3}{(x_3-x_1)(x_3-x_2)(x_3-x_4)}\\ &+\frac{x_4^2\ln x_4}{(x_4-x_1)(x_4-x_2)(x_4-x_3)}\Big].\end{aligned}\tag{E.8}$$

References

[1] For reviews see, for example, H. P. Nilles, Phys. Rep. **110**(1984)1; H. Dreiner, hep-ph/9707435.

[2] J. Erler, J. Feng, N. Polonsky, Phys. Rev. Lett. **78**(1997)3012; J. Ellis, G. Gelmini, C. Jarlskog, G. G. Ross, J. W. F. Valle, Phys. Lett. B**151**(1985)375; R. Barbieri, A. Masiero, Nucl. Phys. B**267**(1986)679; S. Dimopoulos, L. J. Hall, Phys. Lett. B**207**(1987)210; S. Roy, B. Mukhopadhyaya, Phys. Rev. D**55**(1997)7020; H. P. Nilles, N. Polonsky, Nucl. Phys. B**484**(1997)33; R. Hempfling, Nucl. Phys. B**478**(1996)3; M. Hirsch, J. W. F. Valle, Nucl. Phys. B**557**(1999)60; F. de Campos, M.A. Diaz, O.J.P. Eboli, M.B. Ma- gro, L. Navarro, W. Porod, D.A. Restrepo, J.W.F. Valle, hep-ph/9903245.

[3] L. Hall, M. Suzuki, Nucl. Phys. B**231**(1984)419; I. H. Lee, Nucl. Phys. B**246**(1984)120; S. Dawson, Nucl. Phys. B**261**(1985)297;

M. A. Díaz, J. C. Romão, J. W. F. Valle, Nucl. Phys. B**524**(1998)23; C.-H. Chang, T.-F. Feng, Eur. Phys. J. C**12**(2000)137;

[4] D. E. López-Fogliani and C. Muñoz, Phys. Rev. Lett. **97**(2006)041801; N. Escudero, D. E. López-Fogliani, C. Muñoz, and R. Ruiz de Austri, JHEP **0812**(2008)099; J. Fidalgo, D. E. López-Fogliani, C. Muñoz, and R. Ruiz de Austri, JHEP **1110**(2011)020.

[5] J. E. Kim and H. P. Nilles, Phys. Lett. B**138**(1984)150.

[6] For reviews, see H. P. Nilles, Phys. Rept. **110**(1984)1; H. E. Haber and G. L. Kane, Phys. Rept. **117**(1985)75; H. E. Haber, hep-ph/9306207; S. P. Martin, hep-ph/9709356; J. Rosiek, Phys. Rev. D**41**(1990)3464 [hep-ph/9511250].

[7] T.-F. Feng, X.-Q. Li, Phys. Rev. D**63**(2001)073006 and references therein.

[8] Y. Fukuda *et al.*, [Super Kamiokande Collaboration], Phys. Rev. Lett **81**(1998)1562.

[9] Q. R. Ahmad *et al.*, [SNO Collaboration], Phys. Rev. Lett **37**(2001)071301.

[10] K. Eguchi *et al.*, [Kamland Collaboration], Phys. Rev. Lett **90**(2003)021802.

[11] B. Pontecorvo, Sov. Phys. JETP**7**(1958)172; Zh. Eksp. Teor. Fiz.**34**(1958)247; Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys.**28**(1962)870.

[12] D. V. Forero, M. Tórtola and J. W. F. Valle, Phys. Rev. D**86**(2012)073012, [arXiv:1205.4018].

[13] J. Beringer *et al.*, Phys. Rev. D**86**(2012)010001.

[14] F. P. An *et al.*, Phys. Rev. Lett. **108**(2012)171803.

[15] R. Hempfling, Nucl. Phys. B **478**(1996)3; D.E. Kaplan and A.E. Nelson, JHEP **0001**(2000)033; J. C. Romão, M. A. Díaz, M. Hirsch, W. Porod, and J. W. F. Valle, Phys. Rev. D **61**(2000)071703; M. Hirsch, M. A. Díaz, W. Porod, J. C. Romão, and J. W. F. Valle, Phys. Rev. D **62**(2000)113008.

[16] P. Ghosh and S. Roy, JHEP **0904**(2009)069; P. Ghosh, P. Dey, B. Mukhopadhyaya and S. Roy, JHEP **1005**(2010)087.

[17] A. Bartl, M. Hirsch, S. Liebler, W. Porodc and A. Vicente, JHEP **0905**(2009)120; J. Fidalgo, D. E. López-Fogliani, C. Muñoz, and R. Ruiz de Austri, JHEP **0908**(2009)105.

[18] J. Hisano, T. Moroi, K. Tobe, and M. Yamaguchi, Phys. Rev. D**53**(1996)2442.

[19] T. F. Feng, L. Sun and X. Y. Yang, Nucl. Phys. B**800**(2008)221; T. F. Feng, L. Sun and X. Y. Yang, Phys. Rev. D**77**(2008)116008; T. F. Feng and X. Y. Yang, Nucl. Phys. B**814**(2009)101.

[20] S. M. Bilenky, J. Hosek, and S. T. Petcov, Phys. Lett. B**94**(1980)495; J. Schechter and J. W. F. Valle, Phys. Rev. D**23**(1980)2227; M. Doi *et al.*, Phys. Lett. B**102**(1981)323.

[21] O. A. Kiselev *et al.*, [MEG Collaboration], Nucl. Instrum. Meth. A**604**(2009)304.

[22] D. F. Carvalho, M. E. Gómez, and J. C. Romão, Phys. Rev. D**65**(2002)093013.

[23] M. Bona *et al.*, arXiv:0709.0451 [hep-ex].

[24] E821: G.W. Bennett *et al.*, [Muon (g-2) Collaboration], Phys. Rev. D**73**(2006)072003 .

[25] Peter J. Mohr, Barry N. Taylor, David B. Newell, Rev. Mod. Phys. **80**(2008)633 .

[26] G. Aad *et al.*, [ATLAS Collaboration], Phys. Lett. B**716**(2012)1, arXiv:1207.7214 [hep-ex].

[27] S. Chatrchyan *et al.*, [CMS Collaboration], Phys. Lett. B**716**(2012)30, arXiv:1207.7235 [hep-ex].

[28] J. Schechter and J. W. F. Valle, Phys. Rev. D**25**(1982)774; M. Hirsch and J. W. F. Valle, Nucl. Phys. B**557**(1999)60.